

AF-embedding of crossed products of AH-algebras by $\mathbb{Z}$ and asymptotic AF-embedding

Huaxin Lin

Abstract

Let A be a unital AH-algebra and let $\alpha \in \text{Aut}(A)$ be an automorphism. A necessary condition for $A \rtimes_{\alpha} \mathbb{Z}$ being embedded into a unital simple AF-algebra is the existence of a faithful tracial state. If in addition, there is an automorphism κ with $\kappa_{*1} = -\text{id}_{K_1(A)}$ such that $\alpha \circ \kappa$ and $\kappa \circ \alpha$ are asymptotically unitarily equivalent, then $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra. Consequently, in the case that A is a unital AH-algebra (not necessarily simple) with torsion $K_1(A)$, $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if A admits a faithful α -invariant tracial state. We also show that if A is a unital AT-algebra then $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if A admits a faithful α -invariant tracial state. Consequently, for any unital simple AT-algebra A , $A \rtimes_{\alpha} \mathbb{Z}$ can always be embedded into a unital simple AF-algebra.

If X is a compact metric space and $\Lambda : \mathbb{Z}^2 \rightarrow \text{Aut}(C(X))$ is a homomorphism then $C(X) \rtimes_{\Lambda} \mathbb{Z}^2$ can be asymptotically embedded into a unital simple AF-algebra provided that X admits a strictly positive Λ -invariant probability measure. Consequently $C(X) \rtimes_{\Lambda} \mathbb{Z}^2$ is quasidiagonal if X admits a strictly positive Λ -invariant Borel probability measure.

1 Introduction

Quasidiagonality for crossed product C^* -algebras were studied by Dan Voiculescu ([44]), [45] and [46]). Quasidiagonality in C^* -algebras has been studied for a long time in many different point of views (see [6] for more information, also [12], [41], [18], [39], [40], [47], [1], [2], [3], [13], [9] and [10], to name a few). Let X be a compact metric space and α be a homeomorphism on X . It was proved by Pimsner ([37]) that $C(X) \rtimes_{\alpha} \mathbb{Z}$ is quasidiagonal if and only if α is pseudo-non-wondering, and if and only if $C(X) \rtimes_{\alpha} \mathbb{Z}$ can be embedded into an AF-algebra. Recent applications of C^* -algebra theory to dynamical systems further suggest that quasidiagonality and AF-embeddability of crossed products $A \rtimes_{\alpha} \mathbb{Z}$ become increasingly important.

Let A be an AF-algebra and $\alpha \in \text{Aut}(A)$ be an automorphism. Nate Brown ([4]) proved that $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into an AF-algebra if and only if $A \rtimes_{\alpha} \mathbb{Z}$ is quasidiagonal. More importantly, a K -theoretical necessary and sufficient condition for $A \rtimes_{\alpha} \mathbb{Z}$ being quasidiagonal is also given there. When A is a unital simple AT-algebra of real rank zero, Matui ([34]) showed that $A \rtimes_{\alpha} \mathbb{Z}$ can always be embedded into an AF-algebra. On the other hand, Voiculescu's question when $C(X) \rtimes_{\Lambda} \mathbb{Z}^2$ can be embedded into an AF-algebra remains unanswered. It was not even known when $C(X) \rtimes_{\Lambda} \mathbb{Z}^2$ is quasidiagonal in general. It was pointed out by Voiculescu ([46]) that a unital quasidiagonal C^* -algebra always has a tracial state. One consequence of the results here is that if $C(X) \rtimes_{\alpha} \mathbb{Z}^2$ admits a faithful tracial state then it is quasidiagonal (8.2).

We are interested in the problem when C^* -algebras can be embedded into unital simple AF-algebras. Note that not every AF-algebra can be embedded into a unital simple AF-algebra. In fact, it is easy to show that an AF-algebra can be embedded into a unital simple AF-algebra if and only if it admits a faithful tracial state. We prove a more general result. Recall that a C^* -algebra A is AH, if A is an inductive limit of finite direct sums of C^* -algebras A_n with the form

$P_{(n,i)}M_{k(n,i)}(C(X_{n,i}))P_{(n,i)}$, where $X_{n,i}$ is a finite CW complex and $P_{(n,i)} \in M_{k(n,i)}(C(X_{n,i}))$ is a projection. We show that a unital AH-algebra can be embedded into a unital simple AF-algebra if and only if it admits a faithful tracial state (4.1).

Let A be a unital AH-algebra and let $\alpha \in \text{Aut}(A)$ be an automorphism. We study the problem when $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra. A necessary condition is that A admits a faithful α -invariant tracial state. We provide a sufficient condition for $A \rtimes_{\alpha} \mathbb{Z}$ being embedded into a unital simple AF-algebra. We prove this by applying our recent result of so-called Basic Homotopy Lemma in [33]. We show that, when there exists an automorphism κ with $\kappa_{*1} = -\text{id}_{K_1(A)}$ such that $\kappa \circ \alpha$ and $\alpha \circ \kappa$ are asymptotically unitarily equivalent, $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if A admits a faithful α -invariant tracial state (5.8). It should be noted that such κ do exist in many known cases. It also shows that, in the case that A is a unital AH-algebra (not necessarily simple) with torsion $K_1(A)$, $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if A admits a faithful α -invariant tracial state. It should be noted that a unital AH-algebra A with zero $K_1(A)$ may not be AF-algebra. Even in the simple case, a unital simple AH-algebra with zero $K_1(A)$ and with torsion free $K_0(A)$ may not be AF-algebra. There are simple AH-algebras with stable rank higher than one ([42]). On the other hand, we also show that when A is a unital AT-algebra $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if A admits a faithful α -invariant tracial state. In particular, when A is a unital simple AT-algebra, $A \rtimes_{\alpha} \mathbb{Z}$ can always be embedded into a unital simple AF-algebra, which generalizes a result of Matui ([34]).

A C^* -algebra A is said to be asymptotically embedded into a C^* -algebra B , if there exists a sequence of contractive completely positive linear maps $\{\varphi_n\} : A \rightarrow B$ such that $\lim_{n \rightarrow \infty} \|\varphi_n(ab) - \varphi_n(a)\varphi_n(b)\| = 0$ for all $a, b \in A$ and $\lim_{n \rightarrow \infty} \|\varphi_n(a)\| = \|a\|$. Among other things, we show that $C(X) \rtimes_{\Lambda} \mathbb{Z}^2$ can be asymptotically embedded (in a stronger sense) into a unital simple AF-algebra if (and only if) X admits a strictly positive Λ -invariant Borel probability measure (8.1). Consequently, in the case that X admits a strictly positive Λ -invariant Borel probability measure, $C(X) \rtimes_{\Lambda} \mathbb{Z}^2$ is quasidiagonal.

Acknowledgments The preliminary work of this paper was done when the author was in East China Normal University in the summer 2006 which was partially supported by Shanghai Academic Priority Disciplines. This work is also partially supported by a grant from National Science Foundation.

2 Preliminaries

2.1. Let A be a unital C^* -algebra. Denote by $U(A)$ the unitary group of A and denote by $U_0(A)$ the path connected component of $U(A)$ containing the identity. Denote by $\text{Aut}(A)$ the group of automorphisms on A . If $u \in U(A)$, denote by $\text{ad } u$ the inner automorphism defined by $\text{ad } u(a) = u^*au$ for all $a \in A$.

Let A be a unital C^* -algebra and let $\alpha \in \text{Aut}(A)$. Denote by $A \rtimes_{\alpha} \mathbb{Z}$ the crossed product. In this paper, we will fix a unitary and denote it by $u_{\alpha} \in A \rtimes_{\alpha} \mathbb{Z}$ for which $\text{ad } u_{\alpha}(a) = \alpha(a)$ for all $a \in A$.

Let A be a stably finite C^* -algebra. Denote by $T(A)$ the tracial state space of A and by $\text{Aff}(T(A))$ the normed space of all real affine continuous functions on $T(A)$. Denote by $\rho_A : K_0(A) \rightarrow \text{Aff}(T(A))$ the positive homomorphism induced by $\rho_A([e]) = (\tau \otimes T_k)(e)$, where e is a projection in $A \otimes M_k$ and T_k is the standard trace on M_k , $k = 1, 2, \dots$.

Let A and B be two C^* -algebras and let $h : A \rightarrow B$ be a homomorphism. We use h_{*i} for the

induced homomorphism from $K_i(A)$ into $K_i(B)$, $i = 0, 1$.

A homomorphism h is said to be *full*, if $h(a)$ does not belong to any non-trivial (two-sided closed) ideal of B for any $a \neq 0$.

Let $\{B_n\}$ be a sequence of C^* -algebras. Denote by $l^\infty(\{B_n\})$ the C^* -algebraic product of $\{B_n\}$ and $c_0(\{B_n\})$ the C^* -algebraic direct sum. We also use $q_\infty(\{B_n\})$ for the quotient $l^\infty(\{B_n\})/c_0(\{B_n\})$.

A measure μ on X is said to be *strictly positive* measure, if $\mu(O) > 0$ for any non-empty open subset $O \subset X$.

Definition 2.2. Let A and B be two C^* -algebras. Suppose that $L_1, L_2 : A \rightarrow B$ are two maps, $\epsilon > 0$ and $\mathcal{F} \subset A$, we write

$$L_1 \approx_\epsilon L_2 \text{ on } \mathcal{F},$$

if

$$\|L_1(a) - L_2(a)\| < \epsilon \text{ for all } a \in \mathcal{F}.$$

Suppose that $L : A \rightarrow B$ is a contractive completely positive linear map. Let $\delta > 0$ and $\mathcal{G} \subset A$ be a (finite) subset. We say that L is δ - $\mathcal{G}$ -multiplicative if

$$\|L(ab) - L(a)L(b)\| < \delta \text{ for all } a, b \in \mathcal{G}.$$

Let $\sigma > 0$. We say that L is σ - $\mathcal{G}$ -injective if

$$\|L(a)\| \geq \sigma\|a\| \text{ for all } a \in \mathcal{G}.$$

Let $\mathcal{P} \subset \underline{K}(A)$ be a finite subset. A contractive completely positive linear map $L : A \rightarrow B$ is said to be $(\mathcal{G}, \mathcal{P}, \delta)$ map if L is δ - $\mathcal{G}$ -multiplicative, $[L]_{\mathcal{P}}$ is well defined (see 2.2 of [33], 6.1.1 of [24] and see also 2.4 of [9]).

Definition 2.3. Let A and B be two C^* -algebras. Suppose that $\{\varphi_n\}$ is a sequence of contractive completely positive linear maps from A to B . We say $\{\varphi_n\}$ is an *asymptotic embedding*, if

$$\lim_{n \rightarrow \infty} \|\varphi_n(a)\varphi_n(b) - \varphi_n(ab)\| = 0 \text{ for all } a, b \in A \text{ and}$$

$$\lim_{n \rightarrow \infty} \|\varphi_n(a)\| = \|a\| \text{ for all } a \in A.$$

Definition 2.4. Let A and B be two unital C^* -algebras. Let $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times R_+$ be a map and $\mathcal{G} \subset A_+$ be a subset. Suppose that $\varphi : A \rightarrow B$ is a contractive completely positive linear map. We say φ is T - $\mathcal{G}$ -full, if there are $x_1, x_2, \dots, x_{T(a)_\mathbb{N}} \in A$ such that

$$\max\{\|x_i\| : 1 \leq i \leq T(a)_\mathbb{N}\} \leq T(a)_\mathbb{R} \text{ and } \sum_{i=1}^{T(a)_\mathbb{N}} x_i^* \varphi(a) x_i = 1_B$$

for all $a \in \mathcal{G}$.

Suppose that $\{\varphi_n\}$ is a sequence of contractive completely positive linear maps from A to B such that

$$\lim_{n \rightarrow \infty} \|\varphi_n(ab) - \varphi_n(a)\varphi_n(b)\| = 0 \text{ for all } a, b \in A.$$

We say $\{\varphi_n\}$ is a T -full asymptotic embedding if for any finite subset $\mathcal{G} \subset A_+$, there exists an integer $N > 0$ such that φ_n is T - $\mathcal{G}$ -full for all $n \geq N$. From 3.3, it is clear that such $\{\varphi_n\}$ is indeed an asymptotic embedding.

Definition 2.5. Denote by $\mathcal{N}$ the class of separable amenable C^* -algebras which satisfy the Universal Coefficient Theorem.

Definition 2.6. Denote by $\mathcal{U}$ throughout this paper the universal UHF-algebra $\mathcal{U} = \otimes_{n \geq 1} M_n$.

Let $\{e_{i,j}^{(n)}\}$ be the canonical matrix units for M_n . Let $u_n \in M_n$ be the unitary matrix such that $\text{ad } u_n(e_{i,i}^{(n)}) = e_{i+1,i+1}^{(n)}$ (modulo n). Let $\sigma = \otimes_{n \geq 1} \text{ad } u_n \in \text{Aut}(\mathcal{U})$ be the shift (see for example Example 2.2 of [4]). A fact that we will use in this paper is the following property that σ has: For any integer $k > 0$, any $\epsilon > 0$ and any finite subset $\mathcal{F} \subset \mathcal{U}$, there exist mutually orthogonal projections $e_1, e_2, \dots, e_k \in \mathcal{U}$ such that $\sum_{i=1}^k e_i = 1_{\mathcal{U}}$, $\|xe_i - e_i x\| < \epsilon$ for all $x \in \mathcal{F}$ and $\sigma(e_i) = e_{i+1}$, $i = 1, 2, \dots, k$ ($e_{k+1} = e_1$).

Let A be a unital C^* -algebra and let $\alpha \in \text{Aut}(A)$. In this paper, we say that A has the *cyclic Rokhlin property*, if for any integer $k > 0$, any $\epsilon > 0$ and any finite subset $\mathcal{F} \subset A$, there exist mutually orthogonal projections $e_1, e_2, \dots, e_k \in A$ such that

$$\sum_{i=1}^k e_i = 1_A, \quad (\text{e2.1})$$

$$\|xe_i - e_i x\| < \epsilon \text{ for all } x \in \mathcal{F} \text{ and} \quad (\text{e2.2})$$

$$\|\alpha(e_i) - e_{i+1}\| < \epsilon, \quad i = 1, 2, \dots, k \quad (e_{k+1} = e_1). \quad (\text{e2.3})$$

Suppose that α has the above cyclic Rokhlin property and $w \in U(A)$. Then $\text{ad } w \circ \alpha$ also has the cyclic Rokhlin property (a fact communicated to us by N. C. Phillips). Given $\epsilon > 0$ and finite subset $\mathcal{F} \subset A$ and an integer $k > 0$, defined $\mathcal{F}_1 = \{w\} \cup \mathcal{F}$. Since α has the cyclic Rokhlin property, there are mutually orthogonal projections $\{e_1, e_2, \dots, e_k\}$ such that $\sum_{i=1}^k e_i = 1_A$, $\|\alpha(e_i) - e_{i+1}\| < \epsilon/2$ and $\|xe_i - e_i x\| < \epsilon/2$ for all $x \in \mathcal{F}_1$. Then

$$\|\text{ad } w \circ \alpha(e_i) - e_{i+1}\| \leq \|\text{ad } w \circ \alpha(e_i) - \text{ad } w(e_{i+1})\| + \|\text{ad } w(e_{i+1}) - e_{i+1}\| \quad (\text{e2.4})$$

$$< \epsilon/2 + \epsilon/2 = \epsilon. \quad (\text{e2.5})$$

It follows that $\text{ad } w \circ \alpha$ has the cyclic Rokhlin property.

3 Two Uniqueness Theorems

The following is known.

Lemma 3.1. Let A and B be two C^* -algebras. Suppose that $\varphi_n : A \rightarrow B$ is a sequence of contractive completely positive linear map such that

$$\lim_{n \rightarrow \infty} \|\varphi_n(a)\varphi_n(b) - \varphi_n(ab)\| = 0 \text{ for all } a, b \in A.$$

Then the following are equivalent:

- (i) The sequence $\{\varphi_n\}$ is an asymptotic embedding;
- (ii) There exists $\sigma > 0$ and a dense and increasing sequence of finite subset $\mathcal{G}_n \subset A$ such that each φ_n is σ - $\mathcal{G}_n$ -injective;
- (iii) The induced the map: $\Phi : A \rightarrow q_\infty(\{B\})$ defined by $\pi \circ \{\varphi_n\}$, where $\pi : l^\infty(\{B\}) \rightarrow l^\infty(\{B\}) \rightarrow q_\infty(\{B\})$ is the quotient map, is a monomorphism.

Proof. It is clear that (i) implies (ii).

If (ii) holds, then $\pi \circ \{\varphi_n\} : A \rightarrow q_\infty(\{B_n\})$ is a homomorphism. Moreover,

$$\|\pi \circ (\{\varphi_n\})(a)\| \geq \sigma \|a\| \text{ for all } a \in A.$$

It follows that $\pi \circ (\{\varphi_n\})$ is a monomorphism.

If (iii) holds, then

$$\|\pi \circ (\{\varphi_n\})(a)\| = \|a\| \text{ for all } a \in A.$$

It follows that

$$\lim_{n \rightarrow \infty} \|\varphi_n(a)\| = \|a\| \text{ for all } a \in A.$$

□

Lemma 3.2. *Let A be a unital separable amenable C^* -algebra and let $T : A_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$ be a map. Then there exists an increasing sequence of finite subsets $\{\mathcal{G}_n\} \subset A_+$ with its union being dense in A_+ satisfying the following: If B_n is a sequence of unital C^* -algebra and if $\varphi_n : A \rightarrow B_n$ is a contractive completely positive linear map such that*

$$\lim_{n \rightarrow \infty} \|\varphi_n(ab) - \varphi_n(a)\varphi_n(b)\| = 0$$

for all $a \in A$ and each φ_k is T - $\mathcal{G}_n$ full, then the induced map $\Phi = \pi \circ \{\varphi_n\}$ from A to $q_\infty(\{B_n\})$ is a full map, where $\pi : \prod_{n=1}^\infty (\{B_n\} \rightarrow q_\infty(\{B_n\}))$ is the quotient map.

Proof. First we note that $\Phi : A \rightarrow q_\infty(\{B_n\})$ is a homomorphism.

Let $\{\mathcal{G}'_n\}$ be an increasing sequence of finite subsets of A_+ such that $\cup_{n=1}^\infty \mathcal{G}'_n$ is dense in A_+ . For each integer $n > 0$, define $f_{1/n}^{1-1/n}$ to be a positive continuous function on $[0, \infty)$ such that $0 \leq f_{1/n}^{1-1/n}(t) \leq 1$, $f_{1/n}^{1-1/n}(t) = 1$ if $t \geq 1 - 1/n$, $f_{1/n}^{1-1/n}(t) = 0$ if $t \in [0, 1/n]$ and $f_{1/n}^{1-1/n}(t) = \frac{t-1/n}{1-2/n}$ for $t \in [1/n, 1 - 1/n]$.

Let $\mathcal{G}''_n = \{f_{1/n}^{1-1/n}(g) : g \in A_+ \cap \mathcal{G}'_n\}$. Define $\mathcal{G}_n = \mathcal{G}'_n \cup \mathcal{G}''_n$. Now suppose that $\{\varphi_n\}$ is T - $\mathcal{G}_n$ full.

Write $T(a) = (T(a)_\mathbb{N}, T(a)_\mathbb{R})$ for all $a \in A_+ \setminus \{0\}$. Suppose that $a \in A_+ \cap (\cup_{n=1}^\infty \mathcal{G}_n)$ is a non-zero element. It is easy to see that there are $x_1, x_2, \dots, x_N \in q_\infty(\{B_n\})$ with $\|x_i\| \leq T(a)_\mathbb{R}$, $i = 1, 2, \dots, N$ and $N \leq T(a)_\mathbb{N}$ such that

$$\sum_{i=1}^N x_i^* \Phi(a) x_i = 1_{q_\infty(\{B_n\})}.$$

Thus $\Phi(a)$ does not belong to any proper ideal of $q_\infty(\{B_n\})$.

Suppose that $b \in A_+ \setminus \{0\}$ such that $\Phi(b) \in I$ for some (closed two-sided ideal) I of $q_\infty(\{B_n\})$. Let $J = \{b \in A : \Phi(b) \in I\}$. Then J is a (closed two-sided) ideal of A and for any non-zero $a \in A_+ \cap (\cup_{n=1}^\infty \mathcal{G}_n)$, $a \notin J$. Let $b \in J_+ \setminus \{0\}$ with $\|b\| = 1$. Let $0 < 1/n < 1/16$. There exists $c \in A_+ \cap \mathcal{G}'_n$ with $\|c\| \geq 1 - 1/8n$ such that

$$\|c - b\| < 1/2n.$$

By Lemma 2.7 of [19], there is $x \in A$ such that

$$x^*x = a = f_{1/n}^{1-1/n}(c) \text{ and } xx^* \in \overline{bAb}.$$

Therefore $x^*x \in J$. This contradicts with the fact that $a \notin J$.

□

Proposition 3.3. *Let A be a unital separable amenable C^* -algebra and let B be a unital C^* -algebra. Suppose that $\{\varphi_n\}$ is a sequence of contractive completely positive linear maps from A to B such that*

$$\lim_{n \rightarrow \infty} \|\varphi_n(a)\varphi_n(b) - \varphi_n(ab)\| = 0 \text{ for all } a, b \in A.$$

Then there exists a map $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$ such that $\{\varphi_n\}$ is a T -full asymptotic embedding if and only if $\pi \circ (\{\varphi_n\})$ is full.

Proof. Suppose that $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$ is a map and $\{\varphi_n\}$ is a T -full asymptotic embedding. Then, for any $a \in A_+ \setminus \{0\}$, let $N = T(a)_\mathbb{N}$ and $R = T(a)_\mathbb{R}$. Then there are $x_{n,i} \in A$ with $\|x_{n,i}\| \leq R$, $i = 1, 2, \dots, N$ such that

$$\sum_{i=1}^N x_{n,i}^* \varphi_n(a) x_{n,i} = 1_B.$$

Write $X_i = \{x_{n,i}\} \in l^\infty(\{B\})$ (with $\|X_i\| \leq R$). Then

$$\sum_{i=1}^N \pi(X_i)^* \pi \circ (\{\varphi_n\})(a) \pi(X_i) = 1_{q_\infty(\{B\})}.$$

It follows that $\pi \circ (\{\varphi_n\})$ is full.

Conversely, if $\pi \circ (\{\varphi_n\})$ is full, then for any $a \in A_+ \setminus \{0\}$, there is $N \geq 1$ and $R > 0$, and there are $y_i \in q_\infty(\{B\})$ such that $\|y_i\| \leq R$, $i = 1, 2, \dots, N$, and

$$\sum_{i=1}^N y_i^* \pi \circ (\{\varphi_n(a)\}) y_i = 1_{q_\infty(\{B\})}.$$

There is $X_i = \{x_{n,i}\} \in l^\infty(\{B\})$ such that $\|X_i\| \leq R$ and $\pi(X_i) = y_i$, $i = 1, 2, \dots$. Then,

$$\lim_{n \rightarrow \infty} \left\| \sum_{i=1}^N x_{n,i}^* \varphi_n(a) x_{n,i} - 1_B \right\| = 0$$

for all $a \in A_+ \setminus \{0\}$. Thus, for any $a \in A_+ \setminus \{0\}$, there exists an integer $n_a > 0$ such that

$$\left\| \sum_{i=1}^N x_{n,i}^* \varphi_n(a) x_{n,i} - 1_B \right\| < 1/2$$

for all $n \geq n_a$. One obtains an invertible positive element $b_n \in B_+$ with $\|b_n\| \leq 2R$ for all $n \geq n_a$ such that

$$\sum_{i=1}^R b_n x_{n,i}^* \varphi_n(a) x_{n,i} b_n = 1_B$$

for all $n \geq n_a$.

Define $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$ by $T(a)_\mathbb{N} = N$ and $T(a)_\mathbb{R} = 2R$. Then $\{\varphi_n\}$ is T -full. □

Lemma 3.4. *Let A be a separable unital amenable C^* -algebra in $\mathcal{N}$. For any $\epsilon > 0$, any finite subset $\mathcal{F} \subset A$ and any $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$, there exists a finite subset $\mathcal{G} = \mathcal{G}(\epsilon, \mathcal{F}, A) \subset A$, $\delta = \delta(\epsilon, \mathcal{F}, A) > 0$, and a finite subset $\mathcal{P} = \mathcal{P}(\epsilon, \mathcal{F}, A) \subset K_0(A)$, an integer $N = N(\epsilon, \mathcal{F}, A) > 0$ and $\mathcal{S} = \mathcal{S}(\epsilon, \mathcal{F}, A, T) \subset A_+$ satisfying the following: for any unital AF -algebra F with divisible $K_0(F)$ and any T - $\mathcal{S}$ -full acontractive completely positive linear map $f : A \rightarrow F$ with*

$$\|f(a)f(b) - f(ab)\| < \delta \text{ and } \|f(a)\| \geq 1/2\|a\| \quad (\text{e3.6})$$

for all $a, b \in \mathcal{G}$ and any two unital $\mathcal{G}$ - δ -multiplicative contractive completely positive linear maps $\varphi, \psi : A \rightarrow F$ with

$$[\varphi]|_{\mathcal{P}} = [\psi]|_{\mathcal{P}} \quad (\text{e3.7})$$

there exists a unitaries $u \in M_{N+1}(F)$ such that,

$$\|\text{ad } u \circ \text{diag}(\varphi(a), \overbrace{f(a), f(a), \dots, f(a)}^N) - \text{diag}(\psi(a), \overbrace{f(a), f(a), \dots, f(a)}^N)\| < \epsilon \quad (\text{e 3.8})$$

for all $a \in \mathcal{F}$.

Moreover, the assumption that $K_0(F)$ is divisible can be replaced by $K_1(A)$ is torsion free and

$$[f]|_{\mathcal{P}} = [\varphi]|_{\mathcal{P}}. \quad (\text{e 3.9})$$

Proof. Suppose that the lemma is false. Then there are $\epsilon_0 > 0$ and a finite subset $\mathcal{F}_0 \subset A$ and $T : A_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$ such that there are a sequence of decreasing positive numbers $\delta_n \rightarrow 0$, an increasing sequence of finite subsets $\{\mathcal{G}_n\} \subset A$ whose union is dense in A , an increasing sequence of finite subsets $\{\mathcal{P}_n\} \subset K_0(A)$ whose union is $K_0(A)$, an increasing sequence $\{k(n)\}$ of positive integers for which $\lim_{n \rightarrow \infty} k(n) = \infty$, an increasing sequence of finite subsets $\{S_n\}$ which satisfies the conclusion of 3.2, a sequence $\{F_n\}$ of unital AF C^* -algebras with divisible $K_0(F_n)$, and three sequences of $\mathcal{G}_n$ - δ_n -multiplicative contractive completely positive linear maps $\{\varphi_n\}$, $\{\psi_n\}$ and $\{f_n\}$ from A into F_n for which $\{f_n\}$ is T - S_n full and for which

$$[\varphi_n]|_{\mathcal{P}_n} = [\psi_n]|_{\mathcal{P}_n}, \quad (\text{e 3.10})$$

satisfying:

$$\inf\{\sup\{\|u^* \text{diag}(\varphi_n(a), \overbrace{f_n(a), \dots, f_n(a)}^{k(n)})u - \text{diag}(\psi_n(a), \overbrace{f_n(a), \dots, f_n(a)}^{k(n)})\| : a \in \mathcal{F}_0\} \geq \epsilon_0, \quad (\text{e 3.11})$$

where the infimum is taken over all unitaries in $M_{k(n)+1}(B_n)$.

Let $\pi : l^\infty(\{F_n\}) \rightarrow q_\infty(\{F_n\})$ be the quotient map. Put $\bar{\Phi} = \pi \circ \{\varphi_n\}$, $\bar{\Psi} = \pi \circ \{\psi_n\}$ and $\bar{H} = \pi \circ \{f_n\}$. Then $\bar{\Phi}$, $\bar{\Psi}$ and $\bar{H}$ are homomorphisms. By 3.2, $\bar{H}$ is a full monomorphism from A to $q_\infty(\{F_n\})$. Moreover, by (e 3.10), since each F_n is AF (see Cor. 2.1 of [17]),

$$\bar{\Phi}_{*0} = \bar{\Psi}_{*0}. \quad (\text{e 3.12})$$

Since F_n is AF, $K_1(F_n) = \{0\}$ and $K_0(F_n)$ is torsion free. By Cor. 2.1 of [17], since F_n are AF,

$$K_1(q_\infty(\{F_n\}), \mathbb{Z}/k\mathbb{Z}) = \{0\}, \quad k = 2, 3, \dots \quad (\text{e 3.13})$$

Moreover (by Cor. 2.1 of [17]),

$$K_1(q_\infty(\{F_n\})) = \{0\}. \quad (\text{e 3.14})$$

Since $K_0(F_n)$ is divisible,

$$\prod_n K_0(F_n, \mathbb{Z}/k\mathbb{Z}) / \oplus_n K_0(F_n, \mathbb{Z}/k\mathbb{Z}) = \{0\}, \quad i = 0, 1, k = 2, 3, \dots \quad (\text{e 3.15})$$

By Cor. 2.1 of [17],

$$K_0(q_\infty(\{F_n\}), \mathbb{Z}/k\mathbb{Z}) = \{0\}, \quad i = 0, 1, k = 2, 3, \dots \quad (\text{e 3.16})$$

Since A is assumed to satisfy the UCT, we compute that

$$[\bar{\Phi}] = [\bar{\Psi}] \text{ in } KL(A, q_\infty(\{F_n\})). \quad (\text{e 3.17})$$

If the condition that $K_0(F_n)$ is divisible is replaced by $K_1(A)$ is torsion free and $[f_n]|_{\mathcal{P}_n} = [\varphi_n]|_{\mathcal{P}_n}$, then we should get

$$\bar{H}_{*0} = \bar{\Phi}_{*0} = \bar{\Psi}_{*0} \text{ and } K_0(A, \mathbb{Z}/k\mathbb{Z}) = K_0(A)/kK_0(A), \quad k = 2, 3, \dots \quad (\text{e 3.18})$$

Let $\mathcal{Q} \subset \underline{K}(A)$ be a finite subset. There is an integer $k > 0$ such that

$$\mathcal{Q} \cap K_i(A, \mathbb{Z}/m\mathbb{Z}) = \{0\} \text{ for all } m \geq k, \quad (\text{e 3.19})$$

$i = 0, 1$. Define $\tilde{H} = \overbrace{\bar{H} \oplus \bar{H} \oplus \dots \oplus \bar{H}}^{k!-1}$. By (e 3.18), (e 3.13) and (e 3.14), we compute that

$$[\bar{\Phi} \oplus \tilde{H}]|_{\mathcal{Q} \cap K_i(A, \mathbb{Z}/m\mathbb{Z})} = [\bar{\Psi} \oplus \tilde{H}]|_{\mathcal{Q} \cap K_i(A, \mathbb{Z}/m\mathbb{Z})}, \quad (\text{e 3.20})$$

if $i = 2, 3, \dots, k$. By (e 3.14), we also have that

$$[\bar{\Phi} \oplus \tilde{H}]|_{\mathcal{Q} \cap K_1(A)} = [\bar{\Psi} \oplus \tilde{H}]|_{\mathcal{Q} \cap K_1(A)}. \quad (\text{e 3.21})$$

We conclude that

$$[\bar{\Phi} \oplus \tilde{H}]|_{\mathcal{Q}} = [\bar{\Psi} \oplus \tilde{H}]|_{\mathcal{Q}}. \quad (\text{e 3.22})$$

Using either (e 3.17) or (e 3.22) with sufficiently large $\mathcal{Q}$ (which depends on ϵ_0 and $\mathcal{F}_0$) (and hence large k), and by applying 3.9 of [29] we obtain an integer $K > 0$ and a unitary $U \in M_{2K+1}(M_{k!}(q_\infty(\{F_n\})))$ such that

$$\|U^* \text{diag}(\bar{\Phi}(a) \oplus \tilde{H}(a), \overbrace{\tilde{H}(a), \dots, \tilde{H}(a)}^{2K})U - \text{diag}(\bar{\Psi}(a) \oplus \tilde{H}(a), \overbrace{\tilde{H}(a), \dots, \tilde{H}(a)}^{2K})\| < \frac{\epsilon_0}{4} \quad (\text{e 3.23})$$

for all $a \in \mathcal{F}$. Choose $N = (2K+1)(k!-1)$. It is easy to see that there is a unitary $W = \{w_n\} \in l^\infty(\{B_n\})$ such that $\pi(W) = U$. Therefore, there is $N_0 > 0$ such that, for any $n \geq N_0$,

$$\|u_n^* \text{diag}(\varphi_n(a), \overbrace{f_n(a), \dots, f_n(a)}^N)u - \text{diag}(\psi_n(a), \overbrace{f_n(a), \dots, f_n(a)}^N)\| < \frac{\epsilon_0}{2} \quad (\text{e 3.24})$$

for all $a \in \mathcal{F}_0$.

The above estimate (e 3.24) contradicts with (e 3.11). □

Definition 3.5. Denote by $\mathcal{C}_0$ the class of unital C^* -algebra C with the form $C = M_n(C(X))$, where X is a compact subset of a finite CW complex, n is a positive integer, $P \in M_n(C(X))$ is a projection. Note that X is not assumed to be connected.

Recall that a C^* -algebra is said to be AH, if it is an inductive limit of finite direct sums of C^* -algebras in $\mathcal{C}_0$.

The following follows from the results in [31] immediately.

Theorem 3.6. *Let C be a unital AH-algebra and let A be a unital simple C^* -algebra with tracial rank zero. Suppose that $h_1, h_2 : C \rightarrow A$ are two monomorphisms such that*

$$[h_1] = [h_2] \text{ in } KL(C, A) \text{ and} \quad (\text{e 3.25})$$

$$\tau \circ h_1(f) = \tau \circ h_2(f) \text{ for all } f \in C \text{ and } \tau \in T(A). \quad (\text{e 3.26})$$

Then h_1 and h_2 are approximately unitarily equivalent, i.e., there exists a sequence of unitaries $\{u_n\} \subset A$ such that

$$\lim_{n \rightarrow \infty} \text{ad } u_n \circ h_1(f) = h_2(f) \text{ for all } f \in C. \quad (\text{e 3.27})$$

Proof. First, we note that it follows from Cor. 4.8 of [31] (see also 3.4 of [31]) that the theorem holds for $C \in \mathcal{C}_0$. The case that C is a finite direct sum of C^* -algebras in $\mathcal{C}_0$ then also follows.

Now we assume that $C = \lim_{n \rightarrow \infty} (A_n, \varphi_n)$, where each A_n is a finite direct sum of C^* -algebras in $\mathcal{C}_0$. Since the quotients of C^* -algebras in $\mathcal{C}_0$ are still in $\mathcal{C}_0$, without loss of generality, we may assume that each φ_n is injective. Denote by $\varphi_{n,\infty} : A_n \rightarrow C$ the embedding induced by the inductive limit (note we now assume that each φ_n is injective).

Let $\epsilon > 0$ and let $\mathcal{F} \subset C$ be a finite subset. Without loss of generality, we may assume that $\mathcal{F} \subset \varphi_{n,\infty}(A_n)$. Denote by $\mathcal{F}' \subset A_n$ the finite subset so that $\varphi_{n,\infty}(\mathcal{F}') = \mathcal{F}$.

Now

$$[h_1 \circ \varphi_{n,\infty}] = [h_2 \circ \varphi_{n,\infty}] \text{ in } KL(A_n, A) \text{ and}$$

$$\tau \circ h_1 \circ \varphi_{n,\infty}(f) = \tau \circ h_2 \circ \varphi_{n,\infty}(f)$$

for all $f \in A_n$ and $\tau \in T(A)$. By what we have shown (by applying [31]) for the cases that C^* -algebras are direct sum of C^* -algebras in $\mathcal{C}_0$, we obtain a unitary in A such that

$$\text{ad } u \circ h_1 \circ \varphi_{n,\infty} \approx_\epsilon h_2 \circ \varphi_{n,\infty} \text{ on } \mathcal{F}'.$$

It follows that

$$\text{ad } u \circ h_1 \approx_\epsilon h_2 \text{ on } \mathcal{F}.$$

□

4 AH-algebras

Theorem 4.1. *Let A be a unital AH-algebra. Then A can be embedded into a unital simple AF-algebra if and only if A has a faithful tracial state.*

Moreover, if t is a faithful tracial state, then there is a unital simple AF-algebra B with a unique tracial state τ such that $(K_0(B), K_0(B)_+, [1_B]) = (\mathbb{D}, \mathbb{D}_+, 1)$, where $\mathbb{D}$ is a divisible countable dense additive subgroup of $\mathbb{R}$, $K_0(B) = \rho_B(K_0(B))$ and there is a unital monomorphism $\varphi : A \rightarrow B$ such that

$$\tau \circ \varphi = t.$$

Proof. Let t be a faithful tracial state on A . Write $A = \lim_{n \rightarrow \infty} (A_n, h_n)$, where $A_n = \bigoplus_{i=1}^{j(n)} P_{n,j} M_{d(i,n)}(C(X_{n,i})) P_{n,j}$, where each $d(i,n)$ is a positive integer, $X_{n,i}$ is a compact subset of a finite CW complex and $P_{n,j} \in M_{d(i,n)}(C(X_{n,i}))$ is a projection and each h_n is injective. Moreover, we assume that there are integers $r(n,i), s(n,i) > 0$ and a projection

$$Q_{n,i} \in M_{s(n,i)}(P_{n,j} M_{d(i,n)}(C(X_{n,i})) P_{n,j})$$

such that

$$Q_{n,i} M_{s(n,i)}(P_{n,j} M_{d(i,n)}(C(X_{n,i})) P_{n,j}) Q_{n,i} \cong M_{r(n,i)}(C(X_{n,i})).$$

We extend the monomorphism $h_{n,\infty} : M_{s(n,i)}(P_{n,j} M_{d(i,n)}(C(X_{n,i})) P_{n,j}) \rightarrow M_{s(n,i)}(A)$. To simplify notation, we may identify $Q_{n,i} M_{s(n,i)}(P_{n,j} M_{d(i,n)}(C(X_{n,i})) P_{n,j}) Q_{n,i}$ with $M_{r(n,i)}(C(X_{n,i}))$. Let $e_{11}^{(n,i)}$ be a rank one projection in $M_{r(n,i)}(C(X_{n,i}))$. Consider the monomorphism $h_{n,\infty}^{(i)} : e_{11}^{(n,i)} M_{r(n,i)}(C(X_{n,i})) e_{11}^{(n,i)} \rightarrow h_{n,\infty}(e_{11}^{(n,i)}) M_{s(n,i)}(A) h_{n,\infty}(e_{11}^{(n,i)})$ which is the restriction of $h_{n,\infty}$. Note that

$$C(X_{n,i}) \cong e_{11}^{(n,i)} M_{r(n,i)}(C(X_{n,i})) e_{11}^{(n,i)}.$$

We use t for $t \otimes Tr$, where Tr is the standard trace on $M_{r(n,i)}$. This gives a strictly positive probability Borel measure $\mu_{n,i}$ on $X_{n,i}$. As in 3.2 of [32], one obtains a sequence of partitions $P_m : Y_{n,i}(m, j), j = 1, 2, \dots, c(m)$, which has the following properties:

- (1) $Y_{n,i}(m, j) \cap Y_{n,i}(m, j') = \emptyset$, if $j \neq j'$,
- (2) $\lim_{m \rightarrow \infty} \max\{\text{diam}(Y_{n,i}(m, j)) : j = 1, 2, \dots, c(m)\} = 0$,
- (3) $\text{int}(Y_{n,i}(m, j)) \subset Y_{n,i}(m, j) \subset \overline{Y_{n,i}(m, j)}$ and each $Y_{n,i}(m, j)$ is a Borel set,
- (4) $\mu_{n,i}(\partial(Y_{n,i}(m, j))) = 0$,
- (5) $X_{n,i} = \bigcup_{j=1}^{c(m)} Y_{n,i}(m, j)$.

Moreover, $Y_{n,i}(m, j)$ is a union of finitely many members of $Y_{n,i}(m+1, j')$.

Put $D_{n,i} = \{\mu_{n,i}(Y_{n,i}(m, j)) : m, j\}$. Let $\mathbb{D}$ be the additive subgroup of $\mathbb{R}$ generated by $\mathbb{Q}$ and all $D_{n,i}$. Then $\mathbb{D}$ is a divisible countable dense subgroup of $\mathbb{R}$. Let C be a unital simple AF-algebra with $(K_0(C), K_0(C), [1_C]) = (\mathbb{D}, \mathbb{D}_+, 1)$. Let τ be the unique tracial state on C .

Fix n , let $R(n) = \max\{r(n, i) : i = 1, 2, \dots, d(n)\}$. There are mutually orthogonal projections $p_{n,i} \in M_{R(n)}(C)$ such that

$$\tau(p_{n,i}) = t \circ h_{n,\infty}(e_{11}^{(n,i)}), \quad i = 1, 2, \dots, d(n). \quad (\text{e 4.28})$$

Fix i and m , there are mutually orthogonal projections $\{q_{n,i}(m, j)\} \subset p_{n,i} C p_{n,i}$ such that

$$\sigma(e_{n,i}(m, j)) = \mu_{n,i}(Y_{n,i}(m, j)), \quad j = 1, 2, \dots, c(m). \quad (\text{e 4.29})$$

If, for some finite subset J , $\sum_{j' \in J} Y_{n,i}(m+1, j') = Y_{n,i}(m, j)$, then there are mutually orthogonal projections $e_{n,i}(m+1, j') \in e_{n,i}(m, j) C e_{n,i}(m, j)$ such that

$$\sum_{j' \in J} e_{n,i}(m+1, j') = e_{n,i}(m, j) \quad \text{and} \quad (\text{e 4.30})$$

$$\tau(e_{n,i}(m+1, j')) = \mu_{n,i}(Y_{n,i}(m+1, j')). \quad (\text{e 4.31})$$

Therefore

$$\{e_{n,i}(m, j) : j = 1, 2, \dots, c(m), m = 1, 2, \dots\}$$

is a set of mutually orthogonal projections. Let $C(n, i) \subset p_{n,i} C p_{n,i}$ be the C^* -subalgebra generated by $\{e_{n,i}(m, j) : 1 \leq j \leq c(m), m \in \mathbb{N}\}$.

Define $h_m : C(X_{n,i}) \rightarrow p_{n,i} C p_{n,i}$ by

$$h_m^{(n,i)}(f) = \sum_{j=1}^{c(m)} f(\xi_{n,i,m,j}) e_{n,i}(m, j) \quad (\text{e 4.32})$$

for $f \in C(X_{n,i})$, where $\xi_{n,i,m,j} \in Y_{n,i}(m, j)$ is a point. By (2) above and (e 4.30), for each n and i , $\{h_m^{(n,i)}(f)\}$ is a Cauchy sequence. Define

$$h^{(n,i)}(f) = \lim_{m \rightarrow \infty} h_m^{(n,i)}(f) \quad (\text{e 4.33})$$

for $f \in C(X_{n,i})$. We extend, for each n and i , $h^{(n,i)} : M_{r(n,i)}(C(X_{n,i})) \rightarrow M_{r(n,i)}(p_{n,i} C p_{n,i})$. By (e 4.28), (e 4.29) and (e 4.32) as well as (e 4.33), we obtain a monomorphism $h^{(n)} : A_n \rightarrow M_{R(n)}(C)$ such that

$$\tau \circ h^{(n)}(a) = t \circ h_{n,\infty}(a) \quad (\text{e 4.34})$$

for all $a \in A_n$. Since

$$\tau(h^{(n)}(1_{A_n})) \leq 1 \quad \text{and} \quad \ker \rho_C = \{0\}, \quad (\text{e 4.35})$$

there is a unitary $v_n \in M_{R(n)}(C)$ such that $\text{ad } v_n \circ h^{(n)}$ maps A_n into C such that

$$\tau \circ \text{ad } v_n \circ h^{(n)}(a) = t \circ h_{n,\infty}(a) \quad (\text{e 4.36})$$

for all $a \in A_n$. It follows that, for any $n' > n$,

$$\tau \circ \text{ad } v_{n'} \circ h^{(n')} \circ h_{n,n'}(a) = t \circ h_{n',\infty} \circ h_{n,n'}(a). \quad (\text{e 4.37})$$

This implies $(\ker \rho_C = \{0\})$ that

$$(\text{ad } v_{n'} \circ h^{(n')} \circ h_{n,n'})_{*0} = (\text{ad } v_n \circ h^{(n)})_{*0}. \quad (\text{e 4.38})$$

Since $K_1(C) = \{0\}$, $K_0(C)$ is torsion free and divisible, we have

$$[\text{ad } v_{n'} \circ h^{(n')} \circ h_{n,n'}] = [\text{ad } v_n \circ h^{(n)}] \text{ in } KL(A_n, C). \quad (\text{e 4.39})$$

Combining (e 4.39) and (e 4.37), by applying 3.6 and passing to a subsequence, if necessary, we obtain unitaries $u_n \in C$ such that

$$\text{ad } u_{n+1} \circ \text{ad } v_{n+1} \circ h^{(n+1)} \circ h_n \approx_{1/2^n} \text{ad } v_n \circ h^{(n)} \text{ on } \mathcal{F}_n. \quad (\text{e 4.40})$$

Put $\varphi_n = \text{ad } u_n \circ \text{ad } v_n \circ h^{(n)}$, $n = 1, 2, \dots$. Thus we obtain the following approximate intertwining:

$$\begin{array}{ccccccc} A_1 & \xrightarrow{h_1} & A_2 & \xrightarrow{h_2} & A_3 & \xrightarrow{h_3} \dots & A \\ \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \dots \\ C & \xrightarrow{\text{id}_C} & C & \xrightarrow{\text{id}_C} & C & \xrightarrow{\text{id}_C} \dots & C \end{array}$$

Thus, since each φ_n is injective, we obtain a unital embedding $\varphi : A \rightarrow C$ such that $\tau \circ \varphi(a) = t(a)$ for all $a \in A$. □

Corollary 4.2. *Let A be a unital AH-algebra and let $\alpha \in \text{Aut}(A)$. Suppose that A admits a faithful α -invariant tracial state t . Then there is a unital simple AF-algebra B with a unique tracial state τ and with $(K_0(B), K_0(B)_+, [1_B]) = (\mathbb{D}, \mathbb{D}_+, 1)$, where $\mathbb{D} = \rho_B(K_0(B))$, and there is a unital embedding $\varphi : A \rightarrow B$ such that*

$$\tau \circ \varphi \circ \alpha = \tau \circ \varphi.$$

Furthermore, we may assume that $\mathbb{D}$ is divisible.

Proof. In the proof of 4.1, we may begin with the tracial state t . Thus

$$\tau \circ \varphi \circ \alpha = t \circ \alpha = t = \tau \circ \varphi. \quad \square$$

The following is certainly known.

Corollary 4.3. *Let A be a unital AF-algebra. Then A can be embedded into a unital simple AF-algebra if and only if A admits a faithful tracial state.*

5 Crossed products of AH-algebras by $\mathbb{Z}$

The following is proved in [33]. The constant δ and the finite subset $\mathcal{G}$ can be chosen independent of A and h but they are dependent on a measure distribution. Please see 2.10 of [33] for the definition of $\text{Bott}(h, v)|_{\mathcal{P}}$. See 2.10 and 2.11 of [33] also for the definition of $\text{bott}_1(h, v)$ which is related to the usual bott element $\text{bott}_1(u, v)$ for a pair of unitaries.

Theorem 5.1. ([33]) *Let A be a unital AH-algebra and let B be a unital separable simple C^* -algebra of tracial rank zero. Suppose that $h : A \rightarrow B$ is a unital monomorphism. Then, for any $\epsilon > 0$ and any finite subset $\mathcal{F} \subset A$, there exists a positive number δ , a finite subset $\mathcal{G} \subset A$ and a finite subset $\mathcal{P} \subset \underline{K}(A)$ satisfying the following: suppose that $w, u \in U(B)$ are unitaries such that*

$$\|\text{ad } w \circ h(a)u - u \text{ad } w \circ h(a)\| < \delta \text{ for all } a \in \mathcal{G} \text{ and } \text{Bott}(\text{ad } w \circ h, u)|_{\mathcal{P}} = 0.$$

Then there is a continuous rectifiable path $\{u_t : t \in [0, 1]\} \subset U(B)$ such that

$$u_0 = u, \quad u_1 = 1_B \text{ and}$$

$$\|\text{ad } w \circ h(a)u_t - u_t \text{ad } w \circ h(a)\| < \epsilon \text{ for all } a \in \mathcal{F} \text{ and for all } t \in [0, 1].$$

Moreover

$$\|u_t - u_{t'}\| \leq (2\pi + \epsilon)|t - t'| \text{ for all } t, t' \in [0, 1].$$

In particular,

$$\text{Length}(\{u_t\}) \leq 2\pi + \epsilon.$$

Using the above theorem, we obtain the following:

Lemma 5.2. *Let A be a unital AH-algebra and let B be a unital separable simple C^* -algebra of tracial rank zero. Suppose that $h : A \rightarrow B$ is a unital monomorphism.*

Then, for any $\epsilon > 0$ and any finite subset $\mathcal{F} \subset A$, there exists $\delta > 0$ and a finite subset $\mathcal{G} \subset A$ satisfying the following: If there is a continuous path of unitaries $\{u_t : t \in [0, 1]\}$ such that $u_1 = 1_B$,

$$\|h(b)u_t - u_t h(b)\| < \delta \text{ for all } b \in \mathcal{G} \text{ and } t \in [0, 1],$$

then there is a continuous path of unitaries $\{w_t : t \in [0, 1]\}$ such that

$$w_0 = u_0, \quad w_1 = 1_B,$$

$$\|h(a)w_t - w_t h(a)\| < \epsilon \text{ for all } a \in \mathcal{F} \text{ and for all } t \in [0, 1].$$

Moreover,

$$\|u_t - u_{t'}\| \leq (2\pi + \epsilon)|t - t'| \text{ for all } t, t' \in [0, 1].$$

Proof. Fix ϵ and a finite subset $\mathcal{F} \subset A$. Without loss of generality, we may assume that $\mathcal{F}$ is in the unit ball of A . Let $\delta_1 > 0$ and $\mathcal{G}_1 \subset A$ and $\mathcal{P} \subset \underline{K}(A)$ be finite subsets required in 5.1 for the above ϵ and $\mathcal{F}$. We choose $\delta_2 > 0$ and a finite subset $\mathcal{G}'_2 \subset A \otimes C(S^1)$ so that for any δ_2 - $\mathcal{G}'_2$ -multiplicative contractive completely positive linear map $L_1, L_2 : A \otimes C(S^1) \rightarrow B$, $[L_i]|_{\beta(\mathcal{P})}$ is well defined for $i = 1, 2$ and

$$[L_1]|_{\beta(\mathcal{P})} = [L_2]|_{\beta(\mathcal{P})},$$

provided that

$$L_1 \approx_{\delta_2} L_2 \text{ on } \mathcal{G}'_2.$$

(see 2.4 of [33] and 2.10 of [33] for the definition of Bott map β).

Without loss of generality, by choosing smaller δ_2 , we may assume that $\mathcal{G}'_2 = \mathcal{G}_2 \otimes S$, where $S = \{1_{C(S^1)}, z\}$ and z is the identity function on the unit circle.

We may assume that $\mathcal{F} \cup \mathcal{G}_1 \subset \mathcal{G}_2$ and $\delta_2 < \min\{\epsilon, \delta_1\}$.

Let $\delta_3 > 0$ (in place of ϵ) and $\mathcal{G}_3 \subset A$ (in place of $\mathcal{F}_1$) be a finite subset required by 2.7 of [33] for $\delta_2/2$ (in place of ϵ) and $\mathcal{G}_2$ (in place of $\mathcal{F}_2$).

Let $\delta = \min\{\delta_3, \delta_2\}$ and $\mathcal{G} = \mathcal{G}_3 \cup \mathcal{G}_2$.

We may assume that $\mathcal{G}_3$ is a subset of the unit ball of A .

Now suppose that $\{w_t : t \in [0, 1]\}$ and h are in the lemma for the above δ and $\mathcal{G}$.

It follows from 2.7 of [33] that there is, for each t , an δ_2 - $\mathcal{G}_2$ -multiplicative contractive completely positive linear map $\psi_t : A \otimes C(S^1) \rightarrow B$ such that

$$\|\psi_t(a \otimes 1) - h(a)\| < \delta_2/2 \text{ and } \|\psi_t(a \otimes g) - h(a)g(u_t)\| < \delta_2/2 \quad (\text{e 5.41})$$

for all $a \in \mathcal{G}_2$ and $g \in S$.

There is a partition $0 = t_0 < t_1 < \dots < t_m = 1$ such that

$$\|u_{t_{i-1}} - u_{t_i}\| < \delta_2/2, \quad i = 1, 2, \dots, m. \quad (\text{e 5.42})$$

It follows from (e 5.41) and (e 5.42) that

$$\|\psi_{t_{i-1}}(a \otimes g) - \psi_{t_i}(a \otimes g)\| < \delta_2 \text{ for all } a \in \mathcal{G}_2 \text{ and } g \in S. \quad (\text{e 5.43})$$

Moreover,

$$\|\psi_{t_m}(a \otimes g) - h(a)\| < \delta_2 \text{ for all } a \in \mathcal{G}_2 \text{ and } g \in S. \quad (\text{e 5.44})$$

It follows (from the choice of δ_2 and $\mathcal{G}_2$) that

$$[\psi_{t_m}]_{\beta(\mathcal{P})} = 0. \quad (\text{e 5.45})$$

Therefore, by the choice of δ_2 and $\mathcal{G}_2$, and by (e 5.45),

$$[\psi_{t_0}]_{\beta(\mathcal{P})} = 0. \quad (\text{e 5.46})$$

In other words,

$$\text{Bott}(h, u_0)|_{\mathcal{P}} = 0. \quad (\text{e 5.47})$$

Since $\delta < \delta_1$ and $\mathcal{G}_1 \subset \mathcal{G}_2$, by applying 5.1, we obtain a continuous path of unitaries $\{w_t : t \in [0, 1]\}$ such that

$$u_0 = u_0, \quad u_1 = 1_B \text{ and } \|h(a)u_t - u_t h(a)\| < \epsilon \text{ for all } a \in \mathcal{F} \text{ and } t \in [0, 1].$$

Moreover,

$$\|u_t - u_{t'}\| \leq (2\pi + \epsilon)|t - t'| \text{ for all } t, t' \in [0, 1].$$

□

Definition 5.3. Let A be a unital separable C^* -algebra and let B be a unital C^* -algebra. Suppose that $h : A \rightarrow B$ is a unital monomorphism. We say h satisfies property (H) for a positive number L , if the following holds: For any $\epsilon > 0$ and any finite subset $\mathcal{F} \subset A$, if there is a continuous path of unitaries $\{v_t : t \in [0, 1]\}$ in B with $v(1) = 1$ such that

$$\|v_t \text{ad } w \circ h(a) - \text{ad } w \circ h(a)v_t\| < \delta \text{ for all } a \in \mathcal{G} \text{ and } t \in [0, 1],$$

where $w \in B$ is a unitary, there is a continuous path of unitaries $\{u_t : t \in [0, 1]\}$ such that

$$u_0 = v_0, \quad v_1 = 1 \text{ and } \|u_t \text{ad } w \circ h(a) - \text{ad } w \circ h(a)u_t\| < \epsilon$$

for all $a \in \mathcal{F}$ and all $t \in [0, 1]$. Moreover,

$$\|u_t - u_{t'}\| \leq L|t - t'| \text{ for all } t, t' \in [0, 1].$$

It should be noted that a rectifiable continuous path with length, say $2\pi + \epsilon$, may not have the Lipschitz condition.

The following lemma is taken from [34]. The proof is also the same as in [34]. However, since it has different assumption and slightly different conclusion, for the convenience, we will present the proof here.

Lemma 5.4. (Proposition 1 of [34]) *Let A be a unital separable C^* -algebra, let B be a unital simple separable C^* -algebra of tracial rank zero, let $\varphi : A \rightarrow B$ be a unital embedding, let $\alpha \in \text{Aut}(A)$ and let $\beta_0 \in \text{Aut}(B)$ be automorphisms.*

Suppose that $\varphi \circ \alpha$ has the property (H) for a positive number $L > 0$ and suppose that there is a continuous path $\{v(t) : t \in [0, \infty)\}$ of unitaries in $B \otimes \mathcal{U}$ satisfying the following:

$$\lim_{t \rightarrow \infty} \|\varphi \circ \alpha(a) - \text{ad } v(t) \circ \beta \circ \varphi(a)\| = 0$$

for all $a \in A$, where $\beta = \beta_0 \otimes \sigma$ and $\mathcal{U}$ and σ are defined in 2.6.

Then there are unitaries $w, V_n \in B \otimes \mathcal{U}$ ($n = 1, 2, \dots$) such that

$$\varphi' \circ \alpha = \text{ad } w \circ \beta \circ \varphi',$$

where $\varphi'(a) = \lim_{n \rightarrow \infty} \text{ad } (V_1 V_2 \cdots V_n) \circ \varphi$.

Proof. To simplify notation, we may assume that $B = B \otimes \mathcal{U}$. Note that β is an approximately inner automorphism on B which satisfies the cyclic Rokhlin property of 2.6.

Let $\{\mathcal{F}_n\}$ be an increasing sequence of finite subsets of the unit ball of A such that $\overline{\cup_{n=1}^{\infty} \mathcal{F}_n} = A$. Let $N_1 > 0$ such that

$$\frac{L+1}{N_1-1} < 1/16. \quad (\text{e 5.48})$$

Since $\varphi \circ \alpha$ has the property (H) for the constant $L > 0$ and for $\epsilon = 1/16$ and $\mathcal{F} = \cup_{i=0}^{N_1} \alpha^{-i}(\mathcal{F}_1)$, we let $\eta_1 > 0$ (in place of δ) and $\mathcal{G}_1 \subset A$ (in place of $\mathcal{G}$) be a finite subset which is required in 5.3. We may assume that $\mathcal{F}_1 \subset \mathcal{G}_1$.

Choose $0 < \delta_1 < \min\{\frac{\eta_1}{N_1^2}, \frac{1}{32(N_1^2+N_1+1)}\}$.

By replacing β by $\text{ad} \circ v(t_0) \circ \beta$, we may assume that $v(0) = 1$ and

$$\|\varphi \circ \alpha(a) - \text{ad } v(t) \circ \beta \circ \varphi(a)\| < \frac{\delta_1}{2} \quad (\text{e 5.49})$$

for all $a \in \cup_{i=0}^{N_1-1} \alpha^{-i}(\mathcal{G}_1)$ and $t \in [0, \infty)$. In particular,

$$\|\varphi \circ \alpha(a) - \beta \circ \varphi(a)\| < \delta_1/2 \quad \text{and} \quad (\text{e 5.50})$$

$$\|v(t)\varphi \circ \alpha(a)v(t)^* - \varphi \circ \alpha(a)\| < \delta_1/2 \quad (\text{e 5.51})$$

for all $a \in \cup_{i=0}^{N_1-1} \alpha^{-i}(\mathcal{G}_1)$.

Choose $N_2 > N_1$ such that $\frac{L+1}{N_2-1} < \frac{1}{4 \cdot 4^2}$.

Let $1 > \eta_2 > 0$, $\mathcal{G}_2 \subset A$ be a finite subset corresponding to $1/4^3$ and $\cup_{i=0}^{N_2-1} \alpha^{-i}(\mathcal{F}_2)$ in 5.3 (for $\varphi \circ \alpha$ and $L > 0$).

Choose $\delta_2 > 0$ such that $\delta_2 < \min\{\frac{\eta_2}{4}, \frac{1}{4^2(N_2^2+N_2)}\}$.

There is $t_1 > 0$ such that

$$\|\varphi \circ \alpha(a) - \text{ad } v(t) \circ \beta \circ \varphi(a)\| < \frac{\delta_2}{2} \quad (\text{e 5.52})$$

for all $t > t_1$ and $a \in \cup_{i=1}^{N_2-1} \alpha^{-i}(\mathcal{G}_2)$.

For $t \in [0, t_1]$, we define

$$u_0(t) = 1, \quad u_1(t) = v(t), \quad u_2(t) = v(t)\beta(v(t)) \quad (\text{e 5.53})$$

$$u_3(t) = v(t)\beta(v(t))\beta^2(v(t)), \dots \quad (\text{e 5.54})$$

By (e 5.50),

$$\varphi \circ \alpha \approx_{k\delta_1/2} \beta^k \circ \varphi \circ \alpha^{-k+1} \text{ on } \mathcal{G}_1 \quad (\text{e 5.55})$$

$k = 1, 2, \dots, N_1$. Thus,

$$\|[u_k(t), \varphi \circ \alpha(a)]\| = \|[v(t)\beta(v(t)) \cdots \beta^{k-1}(v(t)), \varphi \circ \alpha(a)]\| \quad (\text{e 5.56})$$

$$= \|v(t)\beta(v(t)) \cdots \beta^{k-1}(v(t))\varphi \circ \alpha(a) - v(t)\varphi \circ \alpha(a)\beta(v(t)) \cdots \beta^{k-1}(v(t))\| \quad (\text{e 5.57})$$

$$+ \|v(t)\varphi \circ \alpha(a)\beta(v(t)) \cdots \beta^{k-1}(v(t)) - \varphi \circ \alpha(a)v(t)\beta(v(t)) \cdots \beta^{k-1}(v(t))\| \quad (\text{e 5.58})$$

$$= \|\beta(v(t)) \cdots \beta^{k-1}(v(t))\varphi \circ \alpha(a) - \varphi \circ \alpha(a)\beta(v(t)) \cdots \beta^{k-1}(v(t))\| \quad (\text{e 5.59})$$

$$+ \|v(t)\varphi \circ \alpha(a) - \varphi \circ \alpha(a)v(t)\| \quad (\text{e 5.60})$$

$$= \|\beta(v(t)) \cdots \beta^{k-1}(v(t))\varphi \circ \alpha(a) - \beta(v(t))\varphi \circ \alpha(a)\beta^2(v(t)) \cdots \beta^{k-1}(v(t))\| \quad (\text{e 5.61})$$

$$+ \|\beta(v(t))\varphi \circ \alpha(a)\beta^2(v(t)) \cdots \beta^{k-1}(v(t)) - \varphi \circ \alpha(a)\beta(v(t)) \cdots \beta^{k-1}(v(t))\| \quad (\text{e 5.62})$$

$$+ \|v(t)\varphi \circ \alpha(a) - \varphi \circ \alpha(a)v(t)\| \quad (\text{e 5.63})$$

$$= \|\beta^2(v(t)) \cdots \beta^{k-1}(v(t))\varphi \circ \alpha(a) - \varphi \circ \alpha(a)\beta^2(v(t)) \cdots \beta^{k-1}(v(t))\| \quad (\text{e 5.64})$$

$$+ \|\beta(v(t))\varphi \circ \alpha(a) - \varphi \circ \alpha(a)\beta(v(t))\| \quad (\text{e 5.65})$$

$$+ \|v(t)\varphi \circ \alpha(a) - \varphi \circ \alpha(a)v(t)\|. \quad (\text{e 5.66})$$

$$(\text{e 5.67})$$

It follows that

$$\|[u_k(t), \varphi \circ \alpha(a)]\| \leq \sum_{j=0}^{k-1} \|\beta^j(u(t))\varphi \circ \alpha(a) - \varphi \circ \alpha(a)\beta^j(u(t))\|. \quad (\text{e 5.68})$$

Thus, for $k \leq N_1$, by (e 5.55) and by (e 5.51)

$$\|[u_k(t), \varphi \circ \alpha(a)]\| \leq \sum_{j=0}^{k-1} \|\beta^j(u(t)), \varphi \circ \alpha(a)\| \quad (\text{e 5.69})$$

$$\leq \sum_{j=0}^{k-1} (\|\beta^j(v(t))\beta^j \circ \varphi \circ \alpha^{-j+1}(a) - \beta^j \circ \varphi \circ \alpha^{-j+1}(a)\beta^j(v(t))\| + j\delta_1) \quad (\text{e 5.70})$$

$$\leq \sum_{j=0}^{k-1} (\|v(t)\varphi \circ \alpha^{-j+1}(a) - \varphi \circ \alpha^{-j+1}(a)v(t)\| + j\delta_1) \quad (\text{e 5.71})$$

$$\leq N_1(\delta_1/2 + (N_1 - 1)\delta_1) < N_1^2\delta_1 < \eta_1 \quad (\text{e 5.72})$$

for all $a \in \mathcal{G}_1$ and $t \in [0, t_1]$.

Applying the property (H) for $L > 0$, we obtain a continuous path of unitaries $\{w(t) : t \in [0, 1]\}$ be such that

$$w(0) = u_{N_1}(t_1), \quad w(1) = 1$$

$$\|w(t) - w(t')\| \leq L|t - t'| \text{ and } \|[w(t), \varphi \circ \alpha(a)]\| < \frac{1}{16} \quad (\text{e 5.73})$$

for all $t, t' \in [0, 1]$ and $a \in \cup_{i=1}^{N_1-1} \alpha^{-i}(\mathcal{F}_1)$.

Let $0 = s_0 < s_1 < \dots < s_{N_1-1} = 1$ such that

$$|s_i - s_{i-1}| = \frac{1}{N_1 - 1}, \quad i = 1, 2, \dots, N_1 - 1. \quad (\text{e 5.74})$$

Put $S = \{u_j(t_1), w(s_j) : j = 0, 1, 2, \dots, N_1 - 1\}$. Since β has the cyclic Rokhlin property (see 2.6), we obtain mutually orthogonal projections $e_1, e_2, \dots, e_{N_1} \in B$ such that $\sum_{i=1}^{N_1} e_i = 1_B$ and

$$\|[e_i, b]\| < \frac{L}{128N_1(N_1 - 1)}, \quad i = 1, 2, \dots, N_1 \quad (\text{e 5.75})$$

for $b \in \cup_{i=0}^{N_1-1} \beta^i(S) \cup \cup_{j=0}^{N_1} \alpha^{-j}(\mathcal{G}_1) \cup \varphi(\cup_{j=0}^{N_1} \alpha^{-j+1}(\mathcal{G}_1))$ and with $e_{N_1+1} = e_1$,

$$\|\beta(e_i) - e_{i+1}\| < \frac{L}{128N_1(N_1 - 1)}, \quad i = 1, 2, \dots, N_1. \quad (\text{e 5.76})$$

Define

$$Z = \sum_{j=1}^{N_1} u_{j-1}(t_1) \beta^{j-1}(w(s_{j-1})) e_j. \quad (\text{e 5.77})$$

Denote $\epsilon' = \frac{L}{128N_1(N_1-1)}$. We estimate that, using (e 5.76) and (e 5.75) and the fact that $e_i e_j = 0$ (for $i \neq j$), as well as (e 5.73) and (e 5.74),

$$Z\beta(Z)^* \approx_{3N_1\epsilon'} \sum_{j=1}^{N_1} e_j u_{j-1}(t_1) \beta^{j-1}(w(s_{j-1})) e_j \beta^{j-1}(w(s_{j-2})^*) \beta(u_{j-2}(t_1)^*) \quad (\text{e 5.78})$$

$$\approx_{2N_1\epsilon' + \frac{L}{N_1-1}} \sum_{j=1}^{N_1} e_j u_{j-1}(t_1) \beta(u_{j-2}(t_1)^*) e_j \quad (\text{e 5.79})$$

$$\approx_{N_1\epsilon'} \sum_{j=1}^{N_1} v(t_1) e_j = v(t_1). \quad (\text{e 5.80})$$

Similarly, we then estimate that

$$\|Z^*Z - 1_B\| < \frac{L}{32(N_1 - 1)} \quad \text{and} \quad \|ZZ^* - 1_B\| < \frac{L}{32(N_1 - 1)}. \quad (\text{e 5.81})$$

It follows that there is a unitary $V_1 \in B$ such that

$$\|V_1 - Z\| < \frac{L}{16(N_1 - 1)}. \quad (\text{e 5.82})$$

We also estimate that

$$Z\varphi \circ \alpha(a) = \sum_{j=1}^{N_1} u_{j-1}(t_1) \beta^{j-1}(w(s_{j-1})) e_j \varphi \circ \alpha(a) \quad (\text{e 5.83})$$

$$\approx_{3N_1\epsilon' + \frac{N_1\delta_1}{2}} \sum_{j=1}^{N_1} e_j u_{j-1}(t_1) \beta^{j-1}(w(s_{j-1})) \varphi \circ \alpha^{-j+2}(a) e_j \quad (\text{e 5.84})$$

$$\approx_{1/16} \sum_{j=1}^{N_1} e_j u_{j-1}(t_1) \beta^{j-1} \circ \varphi \circ \alpha^{-j+2}(a) \beta^{j-1}(w(s_{j-1})) e_j \quad (\text{e 5.85})$$

$$\approx_{\frac{N_1\delta_1}{2}} \sum_{j=1}^{N_1} e_j u_{j-1}(t_1) \varphi \circ \alpha(a) \beta^{j-1}(w(s_{j-1})) e_j \quad (\text{e 5.86})$$

$$\approx_{\delta_1/2} \sum_{j=1}^{N_1} e_j \varphi \circ \alpha(a) u_{j-1}(t_1) \beta^{j-1}(w(s_{j-1})) e_j \quad (\text{e 5.87})$$

$$\approx_{3N_1\epsilon'} \varphi \circ \alpha(a) Z \quad (\text{e 5.88})$$

for all $a \in \mathcal{F}_1$.

Note that, since $e_{N_1+1} = e_1$,

$$u_0(t_1)w(0)e_1 = u_{N_1}(t_1)e_1 = u_{N_1}(t)\beta^{N_1}(w(t_{N_1-1}))e_{N_1+1}.$$

Since $\{e_i : 1 \leq i \leq N_1\}$ is mutually orthogonal, as above, we also have

$$v(t_1)\beta(Z) = v(t_1) \sum_{j=1}^{N_1} \beta(u_{j-1}(t_1)) \beta^j(w(t_{j-1})) \beta(e_j) \quad (\text{e 5.89})$$

$$\approx_{4N_1\epsilon'} \sum_{j=1}^{N_1} e_{j+1} v(t_1) \beta(u_{j-1}(t_1)) \beta^j(w(t_{j-1})) e_{j+1} \quad (\text{e 5.90})$$

$$\approx_{\frac{L}{N_1-1}} \sum_{j=1}^{N_1} e_{j+1} v(t_1) \beta(u_{j-1}(t_1)) \beta^j(w(t_j)) e_{j+1} \quad (\text{e 5.91})$$

$$\approx_{2N_1\epsilon'} \sum_{j=1}^{N_1} u_j(t_1) \beta^j(w(t_j)) e_{j+1} \quad (\text{e 5.92})$$

$$= Z. \quad (\text{e 5.93})$$

Therefore (by (e 5.82))

$$\|v(t_1)\beta(V_1) - V_1\| < \left(\frac{(4+128+2)L}{128(N_1-1)}\right) + \frac{2L}{16(N_1-1)} < \frac{1}{8}. \quad (\text{e 5.94})$$

It follows from that

$$\|v(t_1) - V_1\beta(V_1)^*\| < 1/8. \quad (\text{e 5.95})$$

By (e 5.83), (e 5.84), (e 5.85), (e 5.86), (e 5.87), (e 5.88) and (e 5.82)

$$\|[V_1, \varphi \circ \alpha(a)]\| < 3N_1\epsilon' + \frac{N_1\delta_1}{2} + \frac{1}{16} + \frac{N_1\delta_1}{2} + \frac{\delta_1}{2} + 3N_1\epsilon' + \frac{L}{8(N_1-1)} \quad (\text{e 5.96})$$

$$< 1/4 \quad (\text{e 5.97})$$

for all $a \in \mathcal{F}_1$.

Define

$$\begin{aligned} z_1 &= V_1^* v(t_1) \beta(V_1), \quad \varphi_1 = \text{ad } V_1 \circ \varphi, \\ \beta_1 &= \text{ad } z_1^* \circ \beta \quad \text{and} \quad u_1(t) = V_1^* v(t_1) v(t + t_1) V_1. \end{aligned}$$

Then

$$\begin{aligned} \text{adu}_1(t) \circ \beta_1 \circ \varphi_1(a) &= u_1(t)^* z_1 (\beta(V_1)^* \beta \circ \varphi(a) \beta(V_1)) z_1^* u_1(t) \\ &= V_1^* v(t + t_1)^* v(t_1)^* V_1 (V_1^* v(t_1) \beta \circ \varphi(a) v(t_1)^* V_1) V_1^* v(t_1) v(t + t_1) V_1 \\ &= V_1^* v(t + t_1)^* \beta \circ \varphi(a) v(t + t_1) V_1 \end{aligned} \tag{e 5.98}$$

for all $a \in A$. It follows that

$$\lim_{t \rightarrow \infty} \text{adu}_1(t) \circ \beta_1 \circ \varphi_1(a) = \lim_{t \rightarrow \infty} \text{ad } (v(t + t_1) V_1) \circ \beta \circ \varphi(a) \tag{e 5.99}$$

$$= \text{ad } V_1 \circ \varphi \circ \alpha(a) = \varphi_1 \circ \alpha(a) \tag{e 5.100}$$

for all $a \in A$.

We also have $u_1(0) = 1$. By (e 5.94),

$$\|z_1 - 1\| = \|V_1^* v(t_1) \beta(V_1) - V_1^* V_1\| < \frac{1}{8}. \tag{e 5.101}$$

By (e 5.52),

$$\begin{aligned} &\|\varphi_1 \circ \alpha(a) - \text{ad } u_1(t) \circ \beta_1 \circ \varphi_1(a)\| \\ &= \|\varphi \circ \alpha(a) - \text{ad } v(t + t_1) \circ \beta \circ \varphi(a)\| < \frac{\delta_2}{2} \end{aligned} \tag{e 5.102}$$

for all $a \in \cup_{i=0}^{N_2-1} (\mathcal{G}_2)$ and all $t \in [0, \infty)$.

With (e 5.99), (e 5.100) and (e 5.102), we will repeat this process. To repeat the process, we also note that, by 2.6, β_1 has the cyclic Rokhlin property in the sense of 2.6.

By repeating the process, we obtain sequences of unitaries $\{V_n\}$ and $\{z_n\}$ of B such that

$$\|[V_n, \varphi \circ \alpha(a)]\| < \frac{1}{4^n} \tag{e 5.103}$$

for all $a \in \mathcal{F}_n$,

$$\|z_n - 1\| < \frac{1}{4^n}, \tag{e 5.104}$$

$n = 1, 2, \dots$ Moreover, with

$$\varphi_n = \text{ad } (V_1 V_2 \cdots V_{n-1} V_n) \circ \varphi \quad \text{and} \quad \beta_n = \text{ad } (z_1^* z_2^* \cdots z_{n-1}^* z_n^*) \circ \beta,$$

$$\varphi_n \circ \alpha \approx_{\frac{\delta_n}{2}} \beta_n \circ \varphi_n \quad \text{on } \mathcal{F}_n, \tag{e 5.105}$$

$n = 1, 2, \dots$ By (e 5.103), there exists a homomorphism $\varphi' : A \rightarrow B$ such that

$$\varphi'(a) = \lim_{n \rightarrow \infty} \varphi_n(a) \quad \text{for all } a \in A.$$

Since each φ_n is a monomorphism, so is φ' . By (e 5.104), we obtain a unitary $w \in B$ such that

$$w = \lim_{n \rightarrow \infty} z_1^* z_2^* \cdots z_{n-1}^* z_n^*. \tag{e 5.106}$$

It follows from (e 5.99) and (e 5.100) that

$$\varphi' \circ \alpha = \text{ad } w \circ \beta \circ \varphi'.$$

□

Corollary 5.5. *Let A be a unital AH-algebra, let $\alpha \in \text{Aut}(A)$ and let $B \in \mathcal{N}$ be a unital separable simple C^* -algebra with tracial rank zero. Suppose that there is a unital monomorphism $h : A \rightarrow B$ such that there exists a continuous path of unitaries $\{U(t) : t \in [0, \infty)\}$ of B satisfying the following*

$$\lim_{t \rightarrow \infty} \text{ad } U(t) \circ h \circ \alpha(a) = h(a) \text{ for all } a \in A.$$

Then $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra C .

Proof. We first note that $A \otimes \mathcal{U}$ also satisfies the conditions in the statement above with α replaced by $\alpha \otimes \sigma$ and B replaced by $B \otimes \mathcal{U}$, and h by the monomorphism h' defined by $h'(a \otimes b) = h(a) \otimes b$ for all $a \in A$ and $b \in \mathcal{U}$. So we may assume that α satisfy the cyclic Rokhlin property. Next we note that, by 5.2, any unital monomorphism from any unital AH-algebra to a unital separable simple C^* -algebra with tracial rank zero has property (H) for $L = 2\pi + 1$. So, in particular, h has the property (H). Let $\tau \in T(B)$ be a tracial state and let $\mathbb{D}$ be the range of τ on $K_0(B)$. There is a unital simple AF-algebra B_1 with a unique tracial state such that $(K_0(B_1), K_0(B_1)_+, [1_{B_1}]) = (\mathbb{D}, \mathbb{D}_+, 1)$. By the classification theorem of [27], B is a unital simple AH-algebra with real rank zero and with no dimension growth. By [15], there is a unital monomorphism $\varphi : B \rightarrow B_1$. Denote by $C = B_1 \otimes \mathcal{U}$ and $\beta = \text{id}_{B_1} \otimes \sigma$. It follows from 5.4 (and using the cyclic Rokhlin property—see Lemma 4.1 of [32]) that there is a unital monomorphism $\psi : A \rtimes_{\alpha} \mathbb{Z} \rightarrow C \rtimes_{\beta} \mathbb{Z}$. Since $(\beta)_{*0} = \text{id}_{K_0(C)}$ and β has the cyclic Rokhlin property, it follows from Theorem 3.4 of [30] that $C \rtimes_{\beta} \mathbb{Z}$ has tracial rank zero. It follows from the classification theorem (see [27] and [26]) that $C \rtimes_{\beta} \mathbb{Z}$ is a unital simple AH-algebra with slow dimension growth and real rank zero. It follows from [15] that $C \rtimes_{\beta} \mathbb{Z}$ can be embedded into a unital simple AF-algebra. It follows that $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra (with a unique tracial state). $\square$

Lemma 5.6. *Let A be a unital separable amenable C^* -algebra. Then for any finite subset $\mathcal{P} \subset K_1(A)$, there exists $\delta > 0$ and a finite subset $\mathcal{G} \subset A$ satisfying the following: For any pair of automorphisms $\alpha_1, \alpha_2 \in \text{Aut}(A)$, any unital homomorphism $h : A \rightarrow B$ (where B is a unital C^* -algebra) and any unitary $u \in U(B)$, if*

$$\|[h \circ \alpha_i(a), u]\| < \delta \text{ for all } a \in \mathcal{G} \text{ and } (\alpha_1)_{*1} = (-\alpha_2)_{*1}, \quad (\text{e 5.107})$$

then

$$\text{bott}_1(h \circ \alpha_1, u)|_{\mathcal{P}} = -\text{bott}_1(h \circ \alpha_2, u)|_{\mathcal{P}}. \quad (\text{e 5.108})$$

(see 2.10 of [33] for the definition of $\text{bott}_1(\varphi, u)$)

Proof. Suppose that there is a unital C^* -algebra B , there is a homomorphism $\bar{H} : A \rightarrow B$ and a unitary $\bar{U} \in B$ such that $[\bar{H}(a), \bar{U}] = 0$ for all $a \in A$.

Let $\bar{\Phi} : A \otimes C(S^1) \rightarrow B$ be the homomorphism defined by $\bar{\Phi}(a \otimes f) = \bar{H}(a)f(\bar{U})$ for all $a \in A$ and $f \in C(S^1)$. Let $\beta : K_1(A) \rightarrow K_0(A \otimes C(S^1))$ be the homomorphism given by the tensor product. Then $\text{bott}_1(\bar{H} \circ \alpha_i, \bar{U}) = \bar{\Phi}_{*0} \circ \beta \circ (\alpha_i)_{*1}$, $i = 1, 2$. It follows that

$$\text{bott}_1(\bar{H} \circ \alpha_1, \bar{U}) = -\text{bott}_1(\bar{H} \circ \alpha_2, \bar{U}). \quad (\text{e 5.109})$$

To see the lemma holds, we assume that it fails. We then obtain a finite subset $\mathcal{P} \subset K_1(A)$, a sequence of unital C^* -algebras B_n , a sequence of decreasing positive numbers $\{\delta_n\}$ with $\lim_{n \rightarrow \infty} \delta_n = 0$, an increasing sequence of finite subsets $\mathcal{G}_n \subset A$ such that $\cup_{n=1}^{\infty} \mathcal{G}_n$ is dense

in A and a sequence of unital homomorphisms $h_n : A \rightarrow B_n$, and a sequence of unitaries $u_n \in B_n$ such that

$$\|[h_n(a), u_n]\| < \delta_n \text{ for all } a \in \mathcal{G}_n \text{ and} \quad (\text{e 5.110})$$

$$\text{bott}_1(h_n \circ \alpha_1, u_n)|_{\mathcal{P}} \neq -\text{bott}_1(h_n \circ \alpha_2, u_n)|_{\mathcal{P}} \quad (\text{e 5.111})$$

(for all sufficiently large n).

Define a map $H : A \rightarrow l^\infty(\{B_n\})$ and a unitary $U = \{u_n\}$. Denote by $\pi : l^\infty(\{B_n\}) \rightarrow q_\infty(\{B_n\})$ the quotient map. Denote by $\bar{H} = \pi \circ H$ and $\bar{U} = \pi(U)$. Then

$$[\bar{H}(a), \bar{U}] = 0 \text{ for all } a \in A.$$

It follows from what has been proved,

$$\text{bott}_1(\bar{H} \circ \alpha_1, \bar{U}) = -\text{bott}_1(\bar{H} \circ \alpha_2, \bar{U}). \quad (\text{e 5.112})$$

Then, for any sufficiently large n ,

$$\text{bott}_1(h_n \circ \alpha_1, u_n)|_{\mathcal{P}} = -\text{bott}_1(h_n \circ \alpha_2, u_n)|_{\mathcal{P}}. \quad (\text{e 5.113})$$

This contradicts with (e 5.111). □

Lemma 5.7. *Let A be a unital AH -algebra and let B be a unital separable simple C^* -algebra of tracial rank zero with $K_1(B) = \{0\}$ and divisible $K_0(B)$. Let $\alpha, \kappa \in \text{Aut}(A)$ be automorphisms with $\kappa_{*1}(x) = -x$ for all $x \in K_1(A)$. Suppose that there is a unital monomorphism $\varphi : A \rightarrow B$ such that*

$$[\varphi \circ \alpha] = [\varphi] \text{ in } KL(A, B) \quad (\text{e 5.114})$$

and

$$\tau \circ \varphi \circ \alpha = \tau \circ \varphi \quad (\text{e 5.115})$$

for all $\tau \in T(B)$. Suppose also that there is a continuous path of unitaries $\{V(t) : t \in [0, \infty)\}$ of B such that

$$\lim_{t \rightarrow \infty} \text{ad } V(t) \circ \varphi \circ \alpha \circ \kappa(a) = \varphi \circ \kappa \circ \alpha(a) \quad (\text{e 5.116})$$

for all $a \in A$.

Then there is a unital monomorphism $\psi : A \rightarrow M_2(B \otimes \mathcal{U})$ and there is a unitary $w \in U(M_2(B \otimes \mathcal{U}))$ such that

$$\psi \circ \alpha = \text{ad } w \circ (\text{id}_B \otimes \sigma) \circ \psi.$$

Proof. To simplify notation, we may assume that $B = B \otimes \mathcal{U}$ and denote $\beta = \text{id}_B \otimes \sigma$. It is well known that β is approximately inner.

It follows from Theorem 3.6, (e 5.114) and (e 5.115) that there exists a sequence of unitaries $\{u_n\} \subset B$ such that

$$\varphi \circ \alpha(a) = \lim_{n \rightarrow \infty} \text{ad } u_n \circ \beta \circ \varphi(a) \quad (\text{e 5.117})$$

for all $a \in A$.

Define $\psi_0, \bar{\psi}_0 : A \rightarrow M_2(B)$ by

$$\psi_0(a) = \begin{pmatrix} \varphi(a) & 0 \\ 0 & \varphi \circ \kappa \end{pmatrix} \quad \text{and} \quad (\text{e 5.118})$$

$$\bar{\psi}_0 = \begin{pmatrix} \varphi \circ \alpha(a) & 0 \\ 0 & \varphi \circ \alpha \circ \kappa(a) \end{pmatrix} \quad (\text{e 5.119})$$

for all $a \in A$. Define $\beta^{(2)} = \beta \otimes \text{id}_{M_2}$.

It follows from (e 5.117), by passing to a subsequence if necessary, with $U_n = \text{diag}(u_n, u_n)$, we may assume that

$$\bar{\psi}_0(a) = \lim_{n \rightarrow \infty} \text{ad } U_n \circ \beta^{(2)} \circ \psi_0(a) \quad (\text{e 5.120})$$

for all $a \in A$.

Let $\mathcal{F}_1 \subset \mathcal{F}_2 \cdots \subset \mathcal{F}_n \subset \cdots$ be an increasing sequence of the unit ball of B such that the union is dense in the unit ball of B .

Let $1 > \epsilon_n > 0$ such that $\sum_{n=1}^{\infty} \epsilon_n < \infty$. For $\psi_0 : A \rightarrow M_2(B)$, there exists $\delta'_n > 0$, a finite subset $\mathcal{G}'_n \subset A$ and a finite subset $\mathcal{P}_n \subset \underline{K}(A)$ required by 5.1 for ϵ_n and $\mathcal{F}_n$. By choosing larger $\mathcal{P}_n$, we may assume that if $x \in \mathcal{P}_n$, $-x \in \mathcal{P}_n$. For each n , choose $0 < \delta_n \leq \delta'_n$ and a finite subset $\mathcal{G}_n \supset \mathcal{G}'_n$ required in 5.6 for $\mathcal{P}_n \cap K_1(A)$.

Without loss of generality, we may assume that $\mathcal{G}_n$ are in the unit ball of A , $\mathcal{F}_n \subset \mathcal{G}_n \subset \mathcal{G}_{n+1}$, $n = 1, 2, \dots$. Put $S = \{1_{C(S^1)}, z\}$, where $z \in C(S^1)$ is the identity function on the unit circle. Here we also assume that for any δ_n - $\mathcal{G}_n \otimes S$ -multiplicative contractive completely positive linear maps $L_1, L_2 : A \otimes C(S^1) \rightarrow B$ (for any unital C^* -algebra B), $[L_i]|_{\mathcal{P}_n}$ is well defined and

$$[L_1]|_{\mathcal{P}_n} = [L_2]|_{\mathcal{P}_n},$$

provided that

$$L_1 \approx_{\delta_n} L_2 \quad \text{on } \mathcal{G}_n.$$

We may assume that

$$\bar{\psi}_0 \approx_{\delta_n/4} \text{ad } U_n \circ \beta^{(2)} \circ \psi_0 \quad \text{on } \mathcal{G}_n. \quad (\text{e 5.121})$$

Put $W(t) = \text{diag}(1, V(t))$. There are $t_n < t_{n+1} \rightarrow \infty$ such that

$$\text{ad } W(t) \circ \bar{\psi}_0 \approx_{\delta_n/4} \psi_0 \circ \alpha \quad \text{on } \mathcal{G}_n \quad (\text{e 5.122})$$

for all $t \geq t_n$, $n = 1, 2, \dots$

Put

$$Z_n = \begin{pmatrix} u_n^* u_{n+1} & 0 \\ 0 & u_n^* V(t_n)^* V(t_{n+1}) u_{n+1} \end{pmatrix}. \quad (\text{e 5.123})$$

and

$$Z_n(t) = \begin{pmatrix} u_n^* u_{n+1} & 0 \\ 0 & u_n^* V(t_n)^* V(t_{n+1} + (t_n - t_{n+1})t) u_{n+1} \end{pmatrix}. \quad (\text{e 5.124})$$

for $t \in [0, 1]$. So $\{Z_n(t) : t \in [0, 1]\}$ is a path of continuous unitaries, $Z_n(0) = Z_n$ and

$$Z_n(1) = \begin{pmatrix} u_n^* u_{n+1} & 0 \\ 0 & u_n^* u_{n+1} \end{pmatrix} = U_n^* U_{n+1}. \quad (\text{e 5.125})$$

Note that

$$\|[\psi_0 \circ \alpha(a), Z_n(t)]\| < \delta_n \text{ for all } a \in \mathcal{G}_n \text{ and } t \in [0, 1]. \quad (\text{e 5.126})$$

It follows that

$$\text{Bott}(\psi_0, Z_n)|_{\mathcal{P}_n} = \text{Bott}(\psi_0, U_n^* U_{n+1})|_{\mathcal{P}_n}. \quad (\text{e 5.127})$$

By the choice of δ_n and $\mathcal{G}_n$, by applying 5.6,

$$\text{bott}_1(\psi_0, Z_n)|_{\mathcal{P}_n} = \text{bott}_1(\psi_0, U_n^* U_{n+1})|_{\mathcal{P}_n} \quad (\text{e 5.128})$$

$$= \text{bott}_1(\varphi \circ \alpha, u_n^* u_{n+1}) + \text{bott}_1(\varphi \circ \kappa \circ \alpha, u_n^* u_{n+1}) \quad (\text{e 5.129})$$

$$= 0. \quad (\text{e 5.130})$$

Since $K_1(B) = \{0\}$ and $K_0(B)$ is divisible, the above implies that

$$\text{Bott}(\psi_0 \circ \alpha, Z_n)|_{\mathcal{P}_n} = 0, \quad n = 1, 2, \dots \quad (\text{e 5.131})$$

It follows 5.1 that there is a rectifiable continuous path of unitaries $\{Z'_n(t) : t \in [0, 1]\}$ such that

$$Z'_n(1) = Z_n, \quad Z'_n(0) = 1 \text{ and } \|[\psi_0(a), Z'_n(t)]\| < \epsilon_n \quad (\text{e 5.132})$$

for all $a \in \mathcal{F}_n$ and $t \in [0, 1]$. Define

$$v(t) = U_n Z'_n(t - n) \text{ for all } t \in [n, n + 1], \quad n = 1, 2, \dots \quad (\text{e 5.133})$$

Note that $v(n) = U_n Z'_n(0) = U_n$, $v(n + 1) = U_n Z'_n(1) = U_{n+1}$ and $V(t)$ is continuous on $[1, \infty)$. We also compute that

$$\lim_{t \rightarrow \infty} \text{ad } v(t) \circ \beta \circ \psi_0(a) = \psi_0 \circ \alpha(a) \quad (\text{e 5.134})$$

for all $a \in A$.

It follows from 5.2 that $\psi_0 \circ \alpha$ has the property (H) for the constant $2\pi + 1$. Thus Lemma 5.4 applies. $\square$

Theorem 5.8. *Let A be a unital AH-algebra (not necessarily simple) and let $\alpha \in \text{Aut}(A)$ be a automorphism. Suppose that there is another automorphism $\kappa \in \text{Aut}(A)$ such that $\kappa(x) = -x$ for all $x \in K_1(A)$ such that there is a continuous path of unitaries $\{U(t) : t \in [0, \infty)\}$ such that*

$$\lim_{t \rightarrow \infty} \text{ad } U(t) \circ \alpha \circ \kappa(a) = \kappa \circ \alpha(a)$$

for all $a \in A$. Then $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if A has a faithful α -invariant tracial state.

Proof. Suppose that $\varphi : A \rtimes_{\alpha} \mathbb{Z} \rightarrow B$ is a unital monomorphism, where B is a unital simple AF-algebra. Let $\tau \in T(B)$. Then $\tau \circ \varphi$ gives a faithful α -invariant tracial state on A .

So now suppose that there is an α -invariant faithful tracial state τ on A .

It follows from 4.2 that there is a unital simple AF-algebra B with a unique tracial state τ and $(K_0(B), K_0(B)_+, [1_B]) = (\mathbb{D}, \mathbb{D}_+, 1)$, where $\mathbb{D}$ is a dense additive subgroup of $\mathbb{R}$ with $\mathbb{D} = \rho_B(K_0(B))$, such that is a unital embedding $\varphi : A \rightarrow B$ with

$$\tau \circ \varphi \circ \alpha = \tau \circ \varphi. \quad (\text{e 5.135})$$

By considering $B \otimes \mathcal{U}$, we may assume that $K_0(B)$ is divisible. It follows from that fact that $K_0(B) = \mathbb{D}$ and (e 5.135),

$$(\varphi \circ \alpha)_{*0} = \varphi_{*0}. \quad (\text{e 5.136})$$

We also have $(\varphi \circ \alpha)_{*1} = \varphi_{*1}$ since $K_1(B) = \{0\}$. Since $K_0(B)$ is divisible, it follows that

$$[\varphi \circ \alpha] = [\varphi] \text{ in } KL(A, B).$$

It follows from 5.7, there is a unital embedding $\psi : A \rightarrow M_2(B \otimes \mathcal{U})$ and a unitary $w \in U(M_2(B \otimes \mathcal{U}))$ such that

$$\varphi \circ \alpha = \text{ad } w \circ (\text{id}_B \otimes \sigma) \circ \varphi. \quad (\text{e 5.137})$$

Denote by $\beta = \text{id}_B \otimes \sigma$ and $C_1 = M_2(B \otimes \mathcal{U})$. Then we have a unital embedding $\psi_1 : A \rtimes_{\alpha} \mathbb{Z} \rightarrow C_1 \rtimes_{\beta} \mathbb{Z}$. As in the proof of 5.5, this implies that $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra. $\square$

It is worth to note that when $K_1(A) = \{0\}$, one can always take $\kappa = \text{id}_A$.

Corollary 5.9. *Let A be a unital AH-algebra and let $\alpha \in \text{Aut}(A)$. Suppose there is $\kappa \in \text{Aut}(A)$ such that $\alpha(x) = -x$ for all $x \in K_1(A)$ and $\alpha \circ \kappa = \kappa \circ \alpha$. Then $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if there is a faithful α -invariant tracial state*

Proof. This immediately follows from 5.8. $\square$

Corollary 5.10. *Let A be a unital AH-algebra and let $\alpha \in \text{Aut}(A)$. Suppose that $\alpha(x) = -x$ for all $x \in K_1(A)$. Then $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if there is a faithful α -invariant tracial state.*

Proof. Let $\kappa = \alpha$. Then $\kappa \circ \alpha = \alpha \circ \kappa$. So 5.9 applies. $\square$

Corollary 5.11. *Let A be a unital AH-algebra (not necessarily simple) with torsion $K_1(A)$. Let $\alpha \in \text{Aut}(A)$. Then $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if A admits a faithful α -invariant tracial state.*

Proof. By embedding $A \rtimes_{\alpha} \mathbb{Z}$ into $A \otimes \mathcal{U} \rtimes_{\alpha \otimes \sigma} \mathbb{Z}$, we may assume that $K_1(A) = \{0\}$. We then choose $\kappa = \text{id}_A$. So, either 5.9 or 5.8 applies. $\square$

Corollary 5.12. *Let A be a unital AF-algebra and let $\alpha \in \text{Aut}(A)$. Then $A \rtimes_{\alpha} \mathbb{Z}$ can be embedded into a unital simple AF-algebra if and only if A admits a faithful α -invariant tracial state.*

Proof. Every AF-algebra is an AH-algebra with trivial $K_1(A)$. $\square$

Remark 5.13. The scheme of the proof of this section is the same as that in [34]. However, we use 5.1 and 5.2 as well as 3.6 and 4.2. Moreover, by introducing κ , we are able to avoid the difficult generalization of results in [21]. One can verify, for example, for unital simple AT-algebra, required κ can be produced. However, in the next section, we will show that if A is a unital AT-algebra and $\alpha \in \text{Aut}(A)$ is an automorphism then $A \rtimes_{\alpha} \mathbb{Z}$ can always be embedded into a unital simple AF-algebra provided it has a faithful α -invariant tracial state.

It probably worth to point out that, even simple AH-algebras with trivial K_1 -groups and torsion free K_0 -groups are not necessary AF (see [22] for a classification for some of those simple AH-algebras). In particular, they may not have real rank zero. There are also simple AH-algebras whose stable rank is higher than one. Some more exotic AH-algebras were known. In general, when C^* -algebras are not assumed to be simple, there is no realistic hope of classification by K -theory at this moment.

6 Crossed products of $A\mathbb{T}$ -algebras by $\mathbb{Z}$

Some versions of the following three facts are known. We collect here as lemmas for the convenience.

Lemma 6.1. *Let A be a unital C^* -algebra and let $\tau \in T(A)$. Suppose that $u, v \in U(A)$ are unitaries such that*

$$\|u - 1\| < 2/3, \quad \|v - 1\| < 2/3 \quad \text{and} \quad \|uv - 1\| < 2/3. \quad (\text{e 6.138})$$

Then

$$\tau(\log(uv)) = \tau(\log(u)) + \tau(\log(v)). \quad (\text{e 6.139})$$

Proof. Put

$$\lambda = \max\{\|u - 1\|, \|v - 1\|, \|uv - 1\|\}.$$

Let

$$F = \{e^{\pi i t} : -\pi + 2\theta \leq t \leq \pi - 2\theta\},$$

where $\theta = \arccos(3\lambda/2)$.

Then F contains spectra of u , v and uv . Take a branch of logarithm $\log : F \rightarrow (-\pi, \pi)$ which is continuous (smooth) on F . Denote

$$b_1 = \frac{1}{2\pi i} \log(u), \quad b_2 = \frac{1}{2\pi i} \log(v) \quad \text{and} \quad b_3 = \frac{1}{2\pi i} \log(uv).$$

Set

$$w(t) = \begin{cases} e^{-6\pi i t b_1} & 0 \leq t \leq 1/3 \\ e^{-2\pi i b_1} e^{-2\pi i (3t-1)b_2} & 1/3 < t \leq 2/3 \\ e^{-2\pi i b_1} e^{-2\pi i b_2} e^{2\pi i (3t-2)b_3} & 2/3 < t \leq 1. \end{cases}$$

Note that, for each $t \in [0, 1]$, $\text{sp}(w(t)) \subset F$. So $\log w(t)$ is smooth and $\log(w(0)) = \log(w(1))$. One then computes

$$\frac{1}{2\pi i} \int_0^1 \tau\left(\frac{d(w(t))}{dt} w(t)^*\right) dt = 0. \quad (\text{e 6.140})$$

However, one also has that

$$\frac{1}{2\pi i} \int_0^1 \tau\left(\frac{d(w(t))}{dt} w(t)^*\right) dt = \frac{1}{2\pi i} \left\{ \int_0^{1/3} -6\pi i \tau(b_1) dt + \right. \quad (\text{e 6.141})$$

$$\left. + \int_{1/3}^{2/3} 6\pi i \tau(b_2) dt + \int_{2/3}^1 6\pi i \tau(b_3) dt \right\} \quad (\text{e 6.142})$$

$$= -\tau(b_1) - \tau(b_2) + \tau(b_3). \quad (\text{e 6.143})$$

□

Lemma 6.2. *Let A be a unital C^* -algebra, let $B \subset A$ be a unital C^* -subalgebra (with $1_B = 1_A$), let $\alpha : B \rightarrow B$ be a unital homomorphism and let $\tau \in T(A)$ be a tracial state such that $\tau(\alpha(a)) = \tau(a)$ for all $a \in B$. Suppose that $b \in B_{s,a}$ with $\|b\| < \frac{\arcsin(1/6)}{\pi}$, $w \in U(B)$ and $u \in U(A)$ such that*

$$\|w \alpha u(\alpha(w^*)) - 1\| < 1/3 \quad \text{and} \quad \|e^{2\pi i b} w \alpha u(\alpha(w^*) e^{-2\pi i \alpha(b)}) - 1\| < 2/3. \quad (\text{e 6.144})$$

Then

$$\tau(\log(e^{2\pi i b} w \alpha u(\alpha((e^{-2\pi i b} w)^*)))) = \tau(\log(w \alpha u(\alpha(w^*)))). \quad (\text{e 6.145})$$

Proof. By applying 6.1, we have

$$\tau(\log(e^{2\pi ib} w \operatorname{ad} u(\alpha(e^{2\pi ib} w)^*))) \quad (\text{e 6.146})$$

$$= \tau(\log(e^{2\pi ib})) + \tau(\log(w \operatorname{ad} u(\alpha(w^*)e^{-2\pi i\alpha(b)}))) \quad (\text{e 6.147})$$

$$= 2\pi i\tau(b) + \tau(\log(w \operatorname{ad} u(\alpha(w^*))) + \tau(\log(\operatorname{ad} u(e^{-2\pi i\alpha(b)}))) \quad (\text{e 6.148})$$

$$= 2\pi i\tau(b) + \tau(\log(w \operatorname{ad} u(\alpha(w^*))) - 2\pi i\tau(\alpha(b)) \quad (\text{e 6.149})$$

$$= \tau(\log(w \operatorname{ad} u(\alpha(w^*))). \quad (\text{e 6.150})$$

□

Lemma 6.3. *For each integer $k > 0$, there exists $\delta > 0$ satisfying the following:*

Let A be a unital C^* -algebra, let $B \subset A$ be a unital C^* -subalgebra (with $1_B = 1_A$), let $\alpha : B \rightarrow A$ be a unital homomorphism and let $\tau \in T(A)$ be a tracial state such that $\tau(\alpha(a)) = a$ for all $a \in B$, and let $z \in U(B)$.

Suppose that there exists a unitary $u \in U(A)$ such that

$$\|\operatorname{ad} u \circ \alpha(z) - z\| < \delta. \quad (\text{e 6.151})$$

Then

$$\tau(\log(z^m \operatorname{ad} u(\alpha(z^{m*})))) = m\tau(\log(z \operatorname{ad} u(\alpha(z^*)))) \quad (\text{e 6.152})$$

for all $|m| \leq k$.

Proof. With sufficiently small δ , (e 6.151) implies that

$$\|z^m \operatorname{ad} u \circ \alpha(z^{m*}) - 1\| < 1/2^m \text{ and } \|\operatorname{ad} u \circ \alpha(z^{m*})z^m - 1\| < 1/2^m \quad (\text{e 6.153})$$

for all $|m| \leq k$. Then, by 6.1,

$$\begin{aligned} \tau(\log(z^m \operatorname{ad} u(\alpha(z^{m*})))) &= \tau(\log(z^{m-1} \operatorname{ad} u(\alpha(z^{m-1})^*) \operatorname{ad} u(\alpha(z^*))z)) \\ &= \tau(\log(z^{m-1} \operatorname{ad} u(\alpha(z^{m-1})^*))) + \tau(\log(\operatorname{ad} u(\alpha(z^*))z)) \\ &= \tau(\log(z^{m-1} \operatorname{ad} u(\alpha(z^{m-1})^*))) + \tau(\log(z \operatorname{ad} u(\alpha(z^*))))). \end{aligned} \quad (\text{e 6.154})$$

The lemma then follows by induction.

□

Let $B = \oplus_{i=1}^N M_{r(i)}(C(X_i))$, where X_i is a compact subset of the unit circle. Let $e_i \in M_{r(i)}(C(X_i))$ be a constant rank one projection. Denote by z_i the identity function on X_i . We identify $C(X_i)$ with $e_i M_{r(i)}(C(X_i)) e_i$ and z_i with the unitary in $e_i M_{r(i)}(C(X_i)) e_i$, $i = 1, 2, \dots, N$. Denote by $z'_i = (1 - e_i) + z_i$. Denote by E_i the identity of $M_{r(i)}$, the i -th summand of B .

Suppose that A is a unital C^* -algebra of stable rank one and $B \subset A$ with $1_B = 1_A$. Let $\alpha : B \rightarrow A$ be a unital monomorphism. Suppose that $u \in U(A)$ with $uE_i = E_i u$, $i = 1, 2, \dots, N$. Suppose that $\tau \in T(A)$ such that

$$\tau \circ \alpha|_B = \tau|_B.$$

Let $Z = \{v_1, v_2, \dots, v_m\}$ be unitaries in B . Then we have the following:

Proposition 6.4. *There exists $\delta > 0$ and a finite subset $\mathcal{G} \subset B$ satisfying the following:*

If

$$\|\operatorname{ad} u \circ \alpha(a) - a\| < \delta \text{ for all } a \in \mathcal{G} \text{ and} \quad (\text{e 6.155})$$

$$\tau(\log(z_i \operatorname{ad} u(\alpha(z'_i)))) = 0, \quad i = 1, 2, \dots, N, \quad (\text{e 6.156})$$

then

$$\tau(\log(v_j \operatorname{ad} u(\alpha(v_j^*)))) = 0 \quad (\text{e 6.157})$$

for all $v_j \in Z$.

Proof. It is easy to see that we may assume that $E_i = 1_A = 1_B$ and B has only one summand, $B = M_r(C(X))$, where X is a compact subset of the unit circle. Let e be a constant rank one projection in B . We now use z for the identity function on X and the corresponding unitary in eBe and denote $z' = (1 - e) + z$.

For each v_j , we assume that $[v_j] = [z^{m_j}]$ in $K_1(B)$ for some integer m_j . There is a continuous path of unitaries $\{v_j(t) : t \in [0, 1]\}$ such that

$$v_j(0) = v_j \text{ and } v_j(1) = z^{m_j}, \quad j = 1, 2, \dots, N.$$

There are $0 = t_0 < t_1 < \dots < t_K = 1$ such that

$$\|v_j(t_s) - v_j(t_{s-1})\| < 1/3, \quad s = 1, 2, \dots, K \quad (\text{e 6.158})$$

for $1 \leq j \leq N$.

Let $\delta_1 > 0$ be in the lemma 6.3 for $k = \max\{|m_j| : j = 1, 2, \dots, N\}$.

Put

$$\delta = \min\{\delta_1, \frac{1}{6^{K+1}}\}.$$

We now assume that $\mathcal{G}$ contains at least $z, z', v_j(t_{s-1})^* v_j(t_s)$ and $v_j(t_s)$, $s = 1, 2, \dots, N$ and $j = 1, 2, \dots, m$. Now suppose that (e 6.155) and (e 6.156) hold. In particular, we have

$$\|\operatorname{ad} u \circ \alpha(v_j(t_s)) - v_j(t_s)\| < 1/6^{K+1}, \quad (\text{e 6.159})$$

$s = 0, 2, \dots, K$ and $j = 1, 2, \dots, N$. Note that there is $h(j, s) \in B_{s,a}$ with $\|h(j, s)\| < \frac{\arcsin(1/6)}{\pi}$ such that $v_j(t_{s-1})^* v_j(t_s) = e^{2\pi i h(j,s)}$.

By (e 6.156), we have

$$\tau(\log(z' \operatorname{ad} u(\alpha(z')^*))) = 0. \quad (\text{e 6.160})$$

It follows from 6.3 that

$$\tau(\log((z')^{m_j} \operatorname{ad} u(\alpha((z')^{m_j})^*))) = 0, \quad j = 1, 2, \dots, N. \quad (\text{e 6.161})$$

By applying 6.2, we obtain

$$\tau(\log(v_j(t_{N-1}) \operatorname{ad} u \alpha(v_j(t_{N-1})))) = 0. \quad (\text{e 6.162})$$

Then the desired conclusion follows the repeated application of 6.2. □

The following Exel's formula actually holds in much great generality.

Lemma 6.5. *There is a positive number δ satisfying the following:*

Let A be a unital AF-algebra. Suppose that $u, v \in U(A)$ are unitaries and suppose that

$$\|uv - vu\| < \delta. \quad (\text{e 6.163})$$

Then $\operatorname{bott}_1(u, v)$ is well defined and

$$\rho_A(\operatorname{bott}_1(u, v))(\tau) = \frac{1}{2\pi i} \tau(\log(vuv^*u^*)) \text{ for all } \tau \in T(A). \quad (\text{e 6.164})$$

Proof. There is $\delta_1 > 0$ such that $\text{bott}_1(u, v)$ is well defined for any pair of unitaries u and v such that $\|uv - vu\| < \delta_1$. Moreover, there are two projections $p, q \in M_2(A)$ such that

$$\text{bott}_1(u, v) = [p] - [q]. \quad (\text{e 6.165})$$

Now we suppose that $\|uv - vu\| < \delta_1/2$. There is $\delta_2 > 0$ such that

$$\text{bott}_1(u', v) = [p] - [q] \quad (\text{e 6.166})$$

for any other unitary $u' \in U(A)$ satisfying $\|u' - u\| < \delta_2$. Similarly, we obtain $\delta_3 > 0$ such that

$$\text{bott}_1(u', v') = [p] - [q] \quad (\text{e 6.167})$$

for any pair of unitaries $u', v' \in U(A)$ satisfying $\|u' - u\| < \delta_3$ and $\|v' - v\| < \delta_3$.

Now choose $\delta = \delta_1/2$.

There are sequences $\{u_n\}$ and $\{v_n\}$ of unitaries in $\cup_{n=1}^{\infty} A_n$ such that

$$\lim_{n \rightarrow \infty} \|u - u_n\| = 0 \quad \text{and} \quad \lim_{n \rightarrow \infty} \|v - v_n\| = 0. \quad (\text{e 6.168})$$

Without loss of generality, we may assume that

$$u_n, v_n \in A_n, \quad \|u - u_n\| < \delta_3 \quad \text{and} \quad \|v - v_n\| < \delta_3, \quad n \geq N_1 \quad (\text{e 6.169})$$

for some sufficiently large N_1 . Then

$$\text{bott}_1(u_n, v_n) = \text{bott}_1(u, v) = [p] - [q] \quad \text{for all } n \geq N_1. \quad (\text{e 6.170})$$

By choosing larger n , without loss of generality, we may assume that $p, q \in M_2(A_n)$. We may write $A_n = \oplus_{k=1}^N M_{r(k)}$, where $r(k) \geq 1$ is an integer. We may also write

$$p = \oplus_{k=1}^N p_k \quad \text{and} \quad q = \oplus_{k=1}^N q_k, \quad (\text{e 6.171})$$

where $p_k, q_k \in M_{r(k)}$ are projections, $k = 1, 2, \dots, N$. Moreover,

$$u_n = \oplus_{k=1}^N u(k) \quad \text{and} \quad v = \oplus_{k=1}^N v(k), \quad (\text{e 6.172})$$

where $u(k), v(k) \in M_{r(k)}$ are unitaries, $k = 1, 2, \dots, N$.

It follows from Lemma 3.1 of [16] that

$$\text{Tr}_k(p_k) - \text{Tr}_k(q_k) = \frac{1}{2\pi i} \text{Tr}_k(\log(v(k)u(k)v(k)^*u(k)^*)), \quad (\text{e 6.173})$$

where Tr_k denotes the standard trace on $M_{r(k)}$. For any tracial state $\tau \in T(A)$, there are non-negative numbers $\lambda_1, \lambda_2, \dots, \lambda_{R(n)}$ with $\sum_{k=1}^{R(n)} \frac{\lambda_k}{r(k)} = 1$ such that

$$\tau|_{A_n} = \sum_{k=1}^{R(n)} \frac{\lambda_k \text{Tr}_k}{r(k)}. \quad (\text{e 6.174})$$

It follows that

$$\rho_A([p] - [q])(\tau) = \frac{1}{2\pi i} \tau(\log(v_n u_n v_n^* u_n^*)) \quad (\text{e 6.175})$$

for each n and all $\tau \in T(A)$. Note the left side remains the same value. Since we choose a continuous branch of $\log$, we conclude that

$$\rho_A(\text{bott}_1(u, v)) = \frac{1}{2\pi i} \tau(\log(vuv^*u^*)). \quad (\text{e 6.176})$$

□

Lemma 6.6. *Let B be a unital AT -algebra and let A be a unital simple AF -algebra with a unique tracial state τ . Suppose that $\alpha \in \text{Aut}(A)$ and suppose that there is a unital monomorphism $\varphi : B \rightarrow A$ such that*

$$\tau \circ \varphi \circ \alpha = \tau \circ \varphi. \quad (\text{e 6.177})$$

Then there is a unital simple AF -algebra C with a unique tracial state and a unital monomorphism $\psi : A \rightarrow C$ such that there exists a continuous path of unitaries $\{U(t) : t \in [0, \infty)\}$ of C such that

$$\lim_{t \rightarrow \infty} \text{ad } U(t) \circ \psi \circ \varphi \circ \alpha(a) = \psi \circ \varphi(a) \quad (\text{e 6.178})$$

for all $a \in B$.

Proof. By replacing A by $A \otimes \mathcal{U}$, we may assume that $K_0(A)$ is divisible. Suppose that $\rho_A(K_0(A)) = \mathbb{D}$, where $\mathbb{D}$ is a countable divisible dense subgroup of $\mathbb{R}$. By embedding A into another simple AF -algebra, if necessary, we may assume that $(K_0(A), K_0(A)_+, [1_A]) = (\mathbb{D}, \mathbb{D}_+, 1)$. In particular, $\ker \rho_A = \{0\}$.

Let $B = \overline{\bigcup_{n=1}^{\infty} B_n}$, where

$$B_n = \bigoplus_{i=1}^{R(n)} M_{r(i)}(C(X_{n,i}))$$

and $X_{n,i}$ is a compact subset of the unit circle.

Let $e^{(n,i)}$ be a constant rank one projection in $M_{r(i)}(C(X_{n,i}))$. We will identify $C(X_{n,i})$ with $e^{(n,i)} M_{r(i)}(C(X_{n,i})) e^{(n,i)}$ whenever it is convenient. Denote by $z_{n,i}$ the identity function on $X_{n,i}$ and will be identified with a unitary in $e^{(n,i)} M_{r(i)}(C(X_{n,i})) e^{(n,i)}$, $i = 1, 2, \dots, R(n)$, $n = 1, 2, \dots$. Denote $z'_{n,i} = (1 - e_{n,i}) + z_{n,i}$. To simplify the notation, we will identify B with $\varphi(B)$. Also denote by $D_n = \bigoplus_{i=1}^{R(n)} M_{r(i)}$ which is also identified with C^* -subalgebra of B_n of constant matrices.

Since $K_1(A) = \{0\}$, both A and B have torsion K -theory, $\ker \rho_A = \{0\}$, by (e 6.177), in addition to (e 6.177), we also have

$$[\varphi \circ \alpha] = [\varphi] \text{ in } KL(B, A). \quad (\text{e 6.179})$$

So combining (e 6.179) with (e 6.177), by 3.6, there exists a sequence of unitaries $\{w_n\} \subset A$ such that

$$\lim_{n \rightarrow \infty} \text{ad } w_n \circ \alpha(b) = b \quad (\text{e 6.180})$$

for all $b \in B$ (note we have omitted the notation φ).

Write $A = \overline{\bigcup_{n=1}^{\infty} A_n}$, where each A_n is a finite dimensional C^* -subalgebra with $1_A \in A_n$ for each n . We may assume that $\{w_n\} \subset \bigcup_{n=1}^{\infty} A_n$. Moreover, we may assume that

$$\text{ad } w_n \circ \alpha|_{D_n} = \text{id}_{D_n}, \quad n = 1, 2, \dots \quad (\text{e 6.181})$$

By passing to a subsequence if necessarily, we may also assume that

$$\|\text{ad } w_{n+k} \circ \alpha(z_{n,i}) - z_{n,i}\| \quad (\text{e 6.182})$$

is sufficiently small for all $k = 1, 2, \dots$ so that

$$b_{n+k,i} = \frac{1}{2\pi i} \log(z_{n,i} w_{n+k}^* \alpha(z_{n,i}) w_{n+k})$$

can be defined.

Let $\mathbb{D}_1$ be a countable subgroup of $\mathbb{R}$ which contains $\mathbb{D}$ and $\{\tau(b_{n+k,i})\}$. Let C be a unital simple AF-algebra with unique tracial state which we again denote it by τ such that $(K_0(C), K_0(C)_+, [1_C]) = (\mathbb{D}_1, (\mathbb{D}_1)_+, 1)$. For the convenience, we may also assume that $A \subset C$ (and $\tau(a)$ remains the same value) and we use $j : B \rightarrow C$ for the embedding.

Let $\{\mathcal{F}_n\} \subset B$ be an increasing sequence of finite subsets whose union is dense in B . Let $\epsilon_n > 0$ (in place of δ), $\mathcal{G}_n$ (in place of $\mathcal{G}$) be a finite subset of B (in place of A) and $\mathcal{P}_n \subset \underline{K}(B)$ be required in 5.1 for $1/n$ (in place of ϵ) and $\mathcal{F}_n$ (in place of $\mathcal{F}$) (and for j in place of h there). Since B is an AT-algebra, $K_i(B)$ is torsion free ($i = 0, 1$). Therefore, by the fact that $K_1(A) = \{0\}$ and $K_0(A)$ is divisible, it suffices to choose $\mathcal{P}_n \subset K_1(B)$. We may assume that $\epsilon_n \downarrow 0$. Without loss of generality, we may also assume that $\mathcal{F}_n \subset B_n$ and $\mathcal{G}_n$ is in the unit ball of B_n . We may further assume that $z_{n,i} \in \mathcal{G}_n$.

For each $n \geq 2$, let $Z_n = \{z'_{n,i}, z'_{n-1,j}, i, j\}$. To simplify the notation, without loss of generality, we may assume that ϵ_n is smaller than δ in 6.4 and $\mathcal{G}_n$ is larger than $\mathcal{G}$ in 6.4 corresponding B_n and Z_n . Put $\gamma_n = \min\{\tau(e_{n,i}) : 1 \leq i \leq R(n)\}$.

There is $x_{n+k,i} \in K_0(C)$ such that

$$\tau(b_{n+k,i}) = \rho_C(x_{n+k,i}), \quad i = 1, 2, \dots, R(n), n, k = 1, 2, \dots \quad (\text{e 6.183})$$

Let $\mathcal{G}_{n,i} \subset C(X_{n,i})$ be a finite subset such that $\mathcal{G}_n \subset \{\oplus_{i=1}^{R(n)}(a_{k,j}) : a_{k,j} \in \mathcal{G}_{n,i}\}$. Note each $a_{k,j}$ may be viewed as a function in $C(X_{n,i})$.

Let $\delta'_n > 0$ be required by Lemma 2 of [34] for $\frac{\epsilon_n}{2R(n)^2}$. In fact that we choose δ'_n even smaller so that in the conclusion of Lemma 2 of [34] we may require that

$$\|[a_{k,j}(z), w]\| < \epsilon_n/2R(n)^2 \quad (\text{e 6.184})$$

for all $a_{k,j} \in \mathcal{G}_{n,i}$ (in the case that $X_{n,i} = S^1$).

Put $\delta_n = \min\{\epsilon_n/2, \gamma_n \cdot \delta'_n, 2\sin(2\pi\gamma_n \cdot \delta'_n)\}$.

We also assume that the value in (e 6.182) is smaller than δ_n .

We will construct a sequence of unitaries $\{u_n\}$ as follows. For each $X_{n,i} \neq S^1$, choose $\delta(n, i) > 0$ so that

$$\text{bott}_1(z_{n,i}, z) = 0 \quad (\text{e 6.185})$$

for any unitary $z \in e_{n,i}Ce_{n,i}$, provided that

$$\|[z_{n,i}, z]\| < \delta(n, i).$$

We then choose $w_{k(n)}$ so that

$$\|\text{ad } w_{k(n)} \circ \alpha(z_{n,i}) - z_{n,i}\| < \min\{\delta_n, \delta(n, i)\} \quad (\text{e 6.186})$$

$$\text{and } \|\text{ad } w_{k(n)} \circ \alpha(a) - a\| < \delta_n \text{ for all } a \in \mathcal{G}_n. \quad (\text{e 6.187})$$

Moreover,

$$\tau(b_{k(n),i}) < \gamma_n \cdot \delta'_n. \quad (\text{e 6.188})$$

It follows from Lemma 2 of [34] that, if $\text{sp}(z_{n,i}) = S^1$, there is a unitary $v_{n,i} \in e^{(n,i)}Ce^{(n,i)}$ such that

$$\|[z_{n,i}, v_{n,i}]\| < \frac{\epsilon_n}{R(n)^2} \text{ and } \text{bott}_1(z_{n,i}, v_{n,i}) = -x_{k(n),i}. \quad (\text{e 6.189})$$

Let $v_{n,i} = e^{(n,i)}$, if $\text{sp}(z_{n,i}) \neq S^1$. By 6.5 and applying 6.1

$$\tau\left(\frac{1}{2\pi i} \log((z_{n,i} v_{n,i}^* w_{k(n)}^* \alpha(z_{n,i}) w_{n+k} v_{n,i}))\right) \quad (\text{e 6.190})$$

$$= \tau\left(\frac{1}{2\pi i} \log(v_{n,i} (z_{n,i} v_{n,i}^* w_{k(n)}^* \alpha(z_{n,i}) w_{k(n)} v_{n,i} v_{n,i}^*))\right) \quad (\text{e 6.191})$$

$$= \tau\left(\frac{1}{2\pi i} \log(v_{n,i} z_{n,i} v_{n,i}^* z_{n,i}^* z_{n,i} w_{k(n)}^* \alpha(z_{n,i}) w_{k(n)})\right) \quad (\text{e 6.192})$$

$$= \tau\left(\log\left(\frac{1}{2\pi} v_{n,i} z_{n,i} v_{n,i}^* z_{n,i}^*\right)\right) + \tau\left(\log\left(\frac{1}{2\pi} z_{n,i} w_{k(n)}^* \alpha(z_{n,i}) w_{k(n)}\right)\right) \quad (\text{e 6.193})$$

$$= \text{bott}_1(z_{n,i}, v_{n,i}) + \tau\left(\log\left(\frac{1}{2\pi} z_{n,i} w_{k(n)}^* \alpha(z_{n,i}) w_{k(n)}\right)\right) = 0. \quad (\text{e 6.194})$$

We then obtain a unitary u_n (with the form of $\oplus_{i=1}^{R(n)} w_{k(n)} \text{diag}(v_{n,i}, \dots, v_{n,i})$) such that

$$\text{ad } u_n|_{D_n} = \text{id}_{D_n}, \quad \|\text{ad } u_n \circ \alpha(z_{n,i}) - z_{n,i}\| < \epsilon_n/2 \quad (\text{e 6.195})$$

$$\tau\left(\frac{1}{2\pi i} \log(z_{n,i} u_n^* \alpha(z_{n,i}^*) u_n)\right) = 0. \quad (\text{e 6.196})$$

$n = 1, 2, \dots$. By (e 6.187) and choice of δ'_n , we may also assume that

$$\|\text{ad } u_n(\alpha(a)) - a\| < \epsilon_n/2 \text{ for all } a \in \mathcal{G}_n. \quad (\text{e 6.197})$$

By applying 6.4, we also have that

$$\tau\left(\frac{1}{2\pi i} \log(z_{n,j} u_{n+1}^* \alpha(z_{n,j}^*) u_{n+1})\right) = 0. \quad (\text{e 6.198})$$

Put

$$b_1 = \frac{1}{2\pi i} \log(\text{ad } u_{n+1}^*(z_{n,i}) \alpha(z_{n,i}^*)) \quad (\text{e 6.199})$$

$$b_2 = \frac{1}{2\pi i} \log(\text{ad } u_n^*(z_{n,i}) \alpha(z_{n,i}^*)) \quad (\text{e 6.200})$$

$$b_3 = \frac{1}{2\pi i} \log(z_{n,i} \text{ad } u_n^* u_{n+1}(z_{n,i}^*)). \quad (\text{e 6.201})$$

Then, by (e 6.196) and (e 6.198),

$$\tau(b_1) = \tau\left(\frac{1}{2\pi i} \log(z_{n,i} \text{ad}(u_{n+1})(\alpha(z_{n,i})))\right) = 0 \text{ and} \quad (\text{e 6.202})$$

$$\tau(b_2) = \tau\left(\log\left(\frac{1}{2\pi i} z_{n,i} \text{ad } u_n(\alpha(z_{n,i}^*))\right)\right) = 0. \quad (\text{e 6.203})$$

But

$$\text{ad } u_{n+1}^*(e^{2\pi i h_3}) = e^{2\pi i b_1} e^{-2\pi i b_2}. \quad (\text{e 6.204})$$

Therefore, by applying (6.1),

$$\tau(h_3) = \tau\left(\frac{1}{2\pi i} \log(\text{ad } u_{n+1}^*(e^{2\pi i h_3}))\right) = \tau\left(\frac{1}{2\pi i} \log(e^{2\pi i b_1} e^{-2\pi i b_2})\right) \quad (\text{e 6.205})$$

$$= \frac{1}{2\pi i} \tau(2\pi i b_1) + \frac{1}{2\pi i} \tau(2\pi i b_2) = 0. \quad (\text{e 6.206})$$

Note also, by (e 6.197),

$$\|[j(a), u_n^* u_{n+1}]\| < \epsilon_n/2 + \epsilon_{n+1}/2 \text{ for all } a \in \mathcal{G}_n \quad (\text{e 6.207})$$

In particular,

$$\| [z_{n,i}, u_n^* u_{n+1}] \| < \epsilon_n/2 + \epsilon_{n+1}/2. \quad (\text{e 6.208})$$

However, by 6.5

$$\tau(\text{bott}_1(u_{n+1}^* u_n, z_{n,i})) = \tau(h_3) = 0. \quad (\text{e 6.209})$$

Since $\ker \rho_C = \{0\}$,

$$\text{bott}_1(u_{n+1}^* u_n, z_{n,i}) = 0.$$

Therefore

$$\text{bott}_1(z_{n,i}, u_n^* u_{n+1}) = 0. \quad (\text{e 6.210})$$

Hence we may assume that

$$\text{bott}_1(j, u_n^* u_{n+1})|_{\mathcal{P}_n} = 0. \quad (\text{e 6.211})$$

Consequently, (also using (e 6.207)), for sufficiently large n (so ϵ_n is sufficiently small), one may apply 5.1 ($K_i(B)$ is torsion free and $K_1(A) = \{0\}$ and $K_0(A)$ is divisible) to obtain a continuous path of unitaries $\{w_n(t) : t \in [0, 1]\}$ such that

$$w_n(0) = 1, \quad w_n(1) = u_n^* u_{n+1} \quad \text{and} \quad (\text{e 6.212})$$

$$\| [j(a), w_n(t)] \| < 1/n \quad (\text{e 6.213})$$

for all $a \in \mathcal{F}_n$ and $t \in [0, 1]$.

Define

$$U(t+n) = u_n w_n(t) \quad \text{for all } t \in [0, 1]. \quad (\text{e 6.214})$$

Then $U(t)$ is continuous on $[0, \infty)$. One then easily checks that

$$\lim_{n \rightarrow \infty} \text{ad } U(t) \circ \alpha(a) = a \quad \text{for all } a \in B.$$

□

Theorem 6.7. *Let B be a unital AT -algebra and let $\alpha \in \text{Aut}(B)$. Then $A \rtimes_\alpha \mathbb{Z}$ can be embedded into a unital simple AF -algebra if and only if A admits a faithful α -invariant tracial state.*

Proof. It suffices to show the “if” part of the statement. By 4.2, there is a unital simple AF -algebra B with a unique tracial state τ and a unital monomorphism $h : A \rightarrow B$ such that

$$\tau \circ h \circ \alpha = \tau \circ h.$$

It follows from 6.6 that there exists a unital simple AF -algebra C , a unital monomorphism $\varphi : A \rightarrow C$ and a continuous path of unitaries $\{U(t) : t \in [0, \infty)\}$ such that

$$\lim_{t \rightarrow \infty} \text{ad } U(t) \circ \varphi \circ \alpha(a) = \varphi(a) \quad \text{for all } a \in A. \quad (\text{e 6.215})$$

It follows from 5.5 that A can be embedded into a unital simple AF -algebra.

□

Corollary 6.8. *Let A be a unital simple AT -algebra and let $\alpha \in \text{Aut}(A)$. Then $A \rtimes_\alpha \mathbb{Z}$ can always be embedded into a unital simple AF -algebra.*

Proof. Since A is simple, A always admits a faithful α -invariant tracial state. □

Remark 6.9. This section is based on [34]. It simplifies and clarifies some arguments used in [34] for the purpose of proving 6.7. It should be noted that a unital simple AT -algebra may not have real rank zero. Theorem 6.7 also applies to non-simple AT -algebras.

7 Asymptotic embeddings

Lemma 7.1. *Let A be a unital C^* -algebra in $\mathcal{N}$, let $\alpha \in \text{Aut}(A)$ and let C be a unital separable simple AF-algebra. Let $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$. Suppose that there is a T -full unital asymptotic embedding $\{\varphi_n\}$ from A to C such that, for any finite subset $\mathcal{P} \subset K_0(A)$,*

$$[\varphi_n]|_{\mathcal{P}} = [\varphi_n \circ \alpha]|_{\mathcal{P}} \quad (\text{e 7.216})$$

for all sufficiently large n . Then there is a unital asymptotic embedding $\{\psi_n\}$ from A to the unital simple AF-algebra $B = C \otimes \mathcal{U}$ and there is a sequence of unitaries $u_n \in B$ such that

$$\lim_{n \rightarrow \infty} \|\psi_n \circ \alpha(a) - \text{ad } u_n \circ \psi_n(a)\| = 0 \quad (\text{e 7.217})$$

for all $a \in A$. If C has a unique tracial state, one can require that B also has a unique tracial state. Moreover, suppose that $A_0 \subset A$ is a unital C^* -subalgebra and for each finite subset $\mathcal{Q} \in \underline{K}(A_0)$,

$$[\varphi_n]|_{[j_0](\mathcal{Q})} = [\varphi_m]|_{[j_0](\mathcal{Q})} \quad (\text{e 7.218})$$

for all $n, m \geq n_0$ for some $n_0 > 0$, where $j_0 : A_0 \rightarrow A$ is the embedding, then we may also require that

$$[\psi_n]|_{[j_0](\mathcal{Q})} = [\psi_m]|_{[j_0](\mathcal{Q})} \quad (\text{e 7.219})$$

for all sufficiently large n and m .

Proof. We may assume that $\varphi_n(1_A) = 1_B$.

Since $\{\varphi_n\}$ is T -full, by 3.3 and 3.1, for any finite subset $\mathcal{G}_0 \subset A$, there exists $n_0 > 0$ such that

$$\|\varphi_n(a)\| \geq (1/2)\|a\| \text{ for all } a \in \mathcal{G}_0 \quad (\text{e 7.220})$$

for all $n \geq n_0$. Fix a finite subset $\mathcal{F} \subset A$ and $\epsilon > 0$. Let $\mathcal{G} \subset A$, $\mathcal{P} \subset \underline{K}(A)$ be finite subsets, $\delta > 0$ and $N > 0$ be integer meet the requirements of 3.4. Fix $\mathcal{Q} \subset \underline{K}(A_0)$. Without loss of generality, we may assume that $[j_0](\mathcal{Q}) \subset \mathcal{P}$.

Let $K = N(N+1)$. Choose $\mathcal{G}' = \cup_{j=0}^N \alpha^j(\mathcal{G})$. Note that $\pi \circ \{\varphi_n \circ \alpha^j\} : A \rightarrow q_\infty(\{C\})$ is full for each j . Therefore there are $T_j : A_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$, $j = 0, 1, 2, \dots$, such that $\{\varphi_n \circ \alpha^j\}$ are T_j -full. Let $\mathcal{S}_j = \mathcal{S}_j(\epsilon, \mathcal{F}, A, T_j)$ be a finite subset required by 3.4, $j = 0, 1, 2, \dots, N$.

Suppose that $\varphi \in \{\varphi_n\}$ such that it is a $(\mathcal{G}', \mathcal{P}, \delta/2)$ -map. By (e 7.220), we may also assume that

$$\|\varphi(a)\| \geq (1/2)\|a\| \text{ for all } a \in \mathcal{G}'. \quad (\text{e 7.221})$$

We may also assume that

$$[\varphi \circ \alpha^j]|_{\mathcal{P}} = [\varphi]|_{\mathcal{P}}, \quad j = 1, 2, \dots, N. \quad (\text{e 7.222})$$

Define

$$N_a = \max_{0 \leq j \leq K} T(\alpha^j(a))_{\mathbb{N}} \text{ and } R_a = \max_{0 \leq j \leq K} T(\alpha^j(a))_{\mathbb{R}}.$$

Define $T_1 : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$ by $T_1(a) = (N_a, R_a)$ for $a \in A_+ \setminus \{0\}$. Thus we can choose $\varphi \in \{\varphi_n\}$ such that $\varphi \circ \alpha^j$ is T_1 - $\mathcal{S}_j$ full for $j = 1, 2, \dots, N$.

Write $C = C \otimes \mathcal{U}$. Define $\psi : A \rightarrow C \otimes M_K$ by

$$\psi(a) = \text{diag}(\Phi(a), \Phi(\alpha(a)), \dots, \Phi(\alpha^N(a))), \quad (\text{e 7.223})$$

for $a \in A$, where

$$\Phi(b) = \varphi(b) \otimes 1_{M_N} = \text{diag}(\overbrace{\varphi(a), \varphi(a), \dots, \varphi(a)}^N) \quad (\text{e 7.224})$$

for $b \in A$. Thus

$$\psi \circ \alpha(a) = \text{diag}(\Phi(\alpha(a)), \Phi(\alpha^2(a)), \dots, \Phi(\alpha^{N+1}(a))) \quad (\text{e 7.225})$$

for $a \in A$. Note that there is a permutation matrix $W \in M_{N+1}$ such that

$$\text{ad } W \circ \psi(a) = \text{diag}(\Phi(\alpha(a)), \Phi(\alpha^2(a)), \dots, \Phi(\alpha^N(a)), \Phi(a)) \quad (\text{e 7.226})$$

for all $a \in A$.

Since each $\varphi \circ \alpha^j$ is $T_1\text{-}\mathcal{S}_j$ full, by counting multiplicity of each $\varphi(\alpha^j(a))$ for $j = 1, 2, \dots, N$, using (e 7.222) and by applying 3.4, we obtain a unitary $U \in C \otimes M_K$ such that

$$\text{ad } U \circ \psi \approx_\epsilon \psi \circ \alpha \text{ on } \mathcal{F} \quad (\text{e 7.227})$$

By embedding M_K unittally into $\mathcal{U}$, we obtain a $(\mathcal{G}, \mathcal{P}, \delta)$ -map from A into $C \otimes \mathcal{U}$.

By (e 7.222) and (e 7.223), we see that, for any $x \in \mathcal{P}$,

$$[\psi](x) = [\varphi](x) \otimes 1_{\mathcal{U}}. \quad (\text{e 7.228})$$

This proves (e 7.219). Moreover,

$$\|\psi(a)\| \geq (1/2)\|a\| \text{ for all } a \in \mathcal{F}.$$

□

Corollary 7.2. *If in addition to the hypothesis in 7.1, there are $\alpha_1, \alpha_2, \dots, \alpha_k \in \text{Aut}(A)$ such that $\alpha_j \circ \alpha_i = \alpha_i \circ \alpha_j$ ($i, j = 1, 2, \dots, k$) and*

$$\tau \circ \varphi_1(a) = \tau \circ \varphi_n(a) = \tau \circ \varphi_n \circ \alpha_j(a) \quad (\text{e 7.229})$$

for all $a \in A$ and for all $\tau \in T(C)$, $n = 1, 2, \dots$ and $j = 1, 2, \dots, k$, where t is the unique tracial state of $\mathcal{U}$, then we may also require (in addition to the conclusion of 7.1) that

$$(\tau \otimes t) \circ \psi_1(a) = (\tau \otimes t) \circ \psi_n(a) = (\tau \otimes t) \circ \psi_n \circ \alpha_j(a) \quad (\text{e 7.230})$$

for all $a \in A$ and for all $\tau \in T(C)$, $n = 1, 2, \dots$ and $j = 1, 2, \dots, k$.

Proof. Note that, by (e 7.223),

$$\psi \circ \alpha_j(a) = \text{diag}(\Phi(\alpha_j), \Phi(\alpha \circ \alpha_j(a)), \dots, \Phi(\alpha^N \circ \alpha_j(a))). \quad (\text{e 7.231})$$

By (e 7.229) (and (e 7.224)), we have

$$(\tau \otimes t') \circ \Phi(\alpha^i \circ \alpha_j(a)) = (\tau \otimes t') \circ \Phi(\alpha_j(a)) = (\tau \otimes t') \circ \Phi(a) \quad (\text{e 7.232})$$

for all $\tau \in T(C)$, where t' is the tracial state on M_K . It is then easy to conclude that (e 7.230) holds.

□

Lemma 7.3. *Let A be a unital separable amenable C^* -algebra, let $\alpha \in \text{Aut}(A)$ and let B be a unital C^* -algebra. Suppose that there exists a sequence of contractive completely positive linear maps $\{\varphi_n\}$ from A to B which satisfies the following:*

- (1) $\lim_{n \rightarrow \infty} \|\varphi_n(ab) - \varphi_n(a)\varphi_n(b)\| = 0$ for all $a, b \in A$,
- (2) *there exists a sequence of unitaries $\{v_n\}$ in B such that*

$$\lim_{n \rightarrow \infty} \|\text{ad } v_n \circ \varphi_n(a) - \varphi_n(\alpha(a))\| = 0 \quad (\text{e 7.233})$$

for all $a \in A$.

Then there exists a sequence of contractive completely positive linear maps $\{\psi_n\}$ from $A \rtimes_\alpha \mathbb{Z}$ to B such that

$$\lim_{n \rightarrow \infty} \|\psi_n(xy) - \psi_n(x)\psi_n(y)\| = 0 \quad (\text{e 7.234})$$

for all $x, y \in A \rtimes_\alpha \mathbb{Z}$,

$$\lim_{n \rightarrow \infty} \|\psi_n(a) - \varphi_n(a)\| = 0 \text{ and } \lim_{n \rightarrow \infty} \|\psi_n(u_\alpha) - v_n\| = 0 \quad (\text{e 7.235})$$

for all $a \in A$.

Proof. Define

$$L_n\left(\sum_{-N \leq k \leq N} a_k u_\alpha^k\right) = \sum_{-N \leq k \leq N} \varphi_n(a_k) v_n^k \quad (\text{e 7.236})$$

for $a_k \in A$. For any $a, b \in A$ and any integers m and k ,

$$L_n(au_\alpha^m bu_\alpha^k) = L_n(a\alpha^{-m}(b)u_\alpha^{k+m}) = \varphi_n(a\alpha^{-m}(b))v_n^{k+m} \text{ and} \quad (\text{e 7.237})$$

$$L_n(au_\alpha^m)L_n(bu_\alpha^k) = \varphi_n(a)v_n^m \varphi_n(b)v_n^k \quad (\text{e 7.238})$$

$$= [\varphi_n(a)v_n^m \varphi_n(b)v_n^{-m}]v_n^{k+m} \quad (\text{e 7.239})$$

However, by (e 7.233) and (e 7.234),

$$\lim_{n \rightarrow \infty} \|\varphi_n(a\alpha^{-m}(b))v_n^{k+m} - [\varphi_n(a)v_n^m \varphi_n(b)v_n^{-m}]v_n^{k+m}\| = 0 \quad (\text{e 7.240})$$

It follows that

$$\lim_{n \rightarrow \infty} \|L_n(au_\alpha^m bu_\alpha^k) - L_n(au_\alpha)L_n(bu_\alpha^k)\| = 0 \quad (\text{e 7.241})$$

Therefore, for any $x, y \in A \rtimes_\alpha \mathbb{Z}$,

$$\lim_{n \rightarrow \infty} \|L_n(xy) - L_n(x)L_n(y)\| = 0. \quad (\text{e 7.242})$$

Define $L : A \rtimes_\alpha \mathbb{Z} \rightarrow l^\infty(\{B\})$ by $L(c) = \{L_n(c)\}$ for $c \in A \rtimes_\alpha \mathbb{Z}$. Then $\pi \circ L : A \rtimes_\alpha \mathbb{Z} \rightarrow q_\infty(\{B\})$ is a homomorphism. Since A is assumed to be amenable, so is $A \rtimes_\alpha \mathbb{Z}$. It follows from [8] that there is a sequence of contractive completely positive linear maps $\psi_n : A \rtimes_\alpha \mathbb{Z} \rightarrow B$ such that

$$\lim_{n \rightarrow \infty} \|L_n(x) - \psi_n(x)\| = 0 \quad (\text{e 7.243})$$

for all $x \in A \rtimes_\alpha \mathbb{Z} \rightarrow B$. This proves the lemma. $\square$

Corollary 7.4. *Let $A \in \mathcal{N}$ be a unital C^* -algebra, let $\alpha \in \text{Aut}(A)$ and let C be a unital simple AF-algebra. Let $T : A_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$ be a map. Suppose that there is a T -full unital asymptotic embedding $\{\varphi_n\}$ from A to C such that, for any finite subset $\mathcal{P} \subset K_0(A)$,*

$$[\varphi_n]|_{\mathcal{P}} = [\varphi_n \circ \alpha]|_{\mathcal{P}}$$

for all sufficiently large n . Then $A \rtimes_{\alpha} \mathbb{Z}$ can be unittally asymptotically embedded into a unital simple AF-algebra

Proof. By embedding C into $C \otimes \mathcal{U}$, we may assume that $C \cong C \otimes \mathcal{U}$. In particular, $K_0(C)$ is divisible. The corollary then follows from 7.1 and 7.3. $\square$

Corollary 7.5. *Let $A \in \mathcal{N}$ be a unital C^* -algebra, let $\alpha \in \text{Aut}(A)$ and let C be a unital simple AF-algebra. Suppose that there is a monomorphism $\varphi : A \rightarrow C$*

$$\varphi_*0 = (\varphi \circ \alpha)_*0.$$

Then there exists a unital simple AF-algebra B , a sequence of unital monomorphisms $\{\varphi_n\}$ and a sequence of unitaries $\{u_n\}$ in B such that

$$\lim_{n \rightarrow \infty} \|\text{ad } u_n \circ \varphi_n(a) - \varphi_n \circ \alpha(a)\| = 0$$

for all $a \in A$.

Moreover, if C has a unique tracial state τ such that

$$\tau \circ \varphi(a) = \tau \circ \varphi(\alpha(a))$$

for all $a \in A$, then there is an asymptotic embedding $\{\psi_n\} : A \rtimes_{\alpha} \mathbb{Z} \rightarrow B$ such that

$$\lim_{n \rightarrow \infty} \|\psi_n \circ j(a) - \varphi_n(a)\| = 0 \text{ for all } a \in A.$$

We may assume that B has a unique tracial state t and

$$\lim_{n \rightarrow \infty} t(\varphi_n(a)) = \lim_{n \rightarrow \infty} t(\varphi_n(\alpha(a))) = \tau(\varphi(a))$$

for all $a \in A$ as well as

$$[\varphi_n] = [\varphi_n \circ \alpha] \text{ in } KL(A, B). \tag{e 7.244}$$

Proof. The proof is evident by replacing φ_n in the proof of 7.1 by the monomorphism φ . We also apply 7.2. $\square$

Proposition 7.6. *Let A be a unital separable amenable C^* -algebra and let B be a unital C^* -algebra. Suppose that $\alpha \in \text{Aut}(A)$ satisfies the cyclic Rokhlin property (2.6). Suppose that there is a sequence $\{\varphi_n\}$ of contractive completely positive linear maps from $A \rtimes_{\alpha} \mathbb{Z}$ to B for which*

$$\lim_{n \rightarrow \infty} \|\varphi_n(ab) - \varphi_n(a)\varphi_n(b)\| = 0 \text{ for all } a, b \in A.$$

Then $\{\varphi_n\}$ is T -full for some $T : (A \rtimes_{\alpha} \mathbb{Z})_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$ if and only if $\{\varphi_n \circ j\}$ is T' -full for some $T' : A_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$, where $j : A \rightarrow A \rtimes_{\alpha} \mathbb{Z}$ is the natural embedding.

Proof. It suffices to show the “if” part. Suppose that, for some $T' : A_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$, $\{\varphi_n \circ j\}$ is T' -full. Let $\pi : l^\infty(\{B\}) \rightarrow q_\infty(\{B\})$ be the quotient map. Put $\bar{\Phi} = \pi \circ \{\varphi_n\}$. Then $\bar{\Phi} : A \rtimes_\alpha \mathbb{Z} \rightarrow q_\infty(\{B\})$ is a homomorphism. It follows from (3.3) that $\bar{\Phi} \circ j$ is full. Let J_1 be a closed two-sided ideal of $q_\infty(\{B\})$. Suppose that

$$J = \{b \in A \rtimes_\alpha \mathbb{Z} : \bar{\Phi}(b) \in J_1\}.$$

Let $I = J \cap j(A)$. Since α is cyclic Rokhlin, by a result of Pasnicu and Phillips (see 4.1 of [32]), J is proper if and only if I is proper. Since $\bar{\Phi}|_A$ is a full, $I = \{0\}$. So is J . This implies that, for any $b \neq 0$, $\bar{\Phi}(b)$ does not belong to any closed two-sided ideal of $q_\infty(\{B\})$. Thus $\bar{\Phi}$ is full. Therefore, by 3.3, there is $T : (A \rtimes_\alpha \mathbb{Z})_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$ such that $\{\varphi_n\}$ is T -full. $\square$

Corollary 7.7. *Let A be a unital separable amenable C^* -algebra and let B be a unital C^* -algebra. Suppose that $\alpha \in \text{Aut}(A)$ satisfies the cyclic Rokhlin property (2.6). Suppose that there is an asymptotic embedding $\{\varphi_n\} : A \rtimes_\alpha \mathbb{Z} \rightarrow B$ and there is a full monomorphism $h : A \rightarrow B$ such that*

$$\lim_{n \rightarrow \infty} \|h(a) - \varphi_n \circ j(a)\| = 0 \quad (\text{e 7.245})$$

for all $a \in A$, where $j : A \rightarrow A \rtimes_\alpha \mathbb{Z}$ is the natural embedding. Then $\{\varphi_n\}$ is T -full for some $T : (A \rtimes_\alpha \mathbb{Z})_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$.

Proof. It follows from (e 7.245) that $\bar{\Phi} \circ j = h$. Thus $\bar{\Phi} \circ j$ is full. $\square$

8 Asymptotic Embeddings for $C(X) \rtimes_\alpha \mathbb{Z}^2$ and quasidiagonality

Theorem 8.1. *Let X be a compact metric space and let $\Lambda : \mathbb{Z}^2 \rightarrow \text{Aut}(C(X))$ be a $\mathbb{Z}^2$ action on X .*

Then the following are equivalent:

- (1) *there exists a strictly positive Λ -invariant Borel probability measure μ on X ,*
- (2) *there exists a unital simple AF-algebra B , there exists a monomorphism $h : C(X) \rightarrow B$ and there exists a T -full asymptotic embedding $\{\varphi_n\} : C(X) \rtimes_\Lambda \mathbb{Z}^2 \rightarrow B$ for some $T : A_+ \setminus \{0\} \rightarrow (\mathbb{N}, \mathbb{R}_+)$ such that*

$$\lim_{n \rightarrow \infty} \|h(f) - \varphi_n \circ j(f)\| = 0 \quad (\text{e 8.246})$$

for all $f \in C(X)$, where $j : C(X) \rightarrow C(X) \rtimes_\Lambda \mathbb{Z}^2$ is the natural embedding.

Proof. Suppose that (2) holds. Without loss of generality, we may assume that $\varphi_n(1_{C(X) \rtimes_\Lambda \mathbb{Z}^2}) = 1_B$. Let $\tau \in T(C)$ be a tracial state. Then $\tau \circ h$ is a strictly positive Borel probability measure. Since $C(X) \rtimes_\Lambda \mathbb{Z}^2$ is separable, there exists a subsequence $\{n_k\}$ and a tracial state t of $C(X) \rtimes_\Lambda \mathbb{Z}^2$ such that

$$t(a) = \lim_{k \rightarrow \infty} \tau \circ \varphi_{n_k}(a)$$

for all $a \in C(X) \rtimes_\Lambda \mathbb{Z}^2$. Therefore $t \circ j$ is Λ -invariant. However, by (e 8.246)

$$t \circ j = \tau \circ j.$$

Thus $\tau \circ j$ is a strictly positive Λ -invariant Borel probability measure on X .

Now suppose that (1) holds.

It follows from Cor 3.3 and Theorem 4.3 of [32] (and its proof) that there is a unital simple AF-algebra B with a unique tracial state τ and a unital embedding $\varphi : C(X) \otimes \mathcal{U} \rtimes_{\gamma_1} \mathbb{Z} \rightarrow B$, such that

$$\tau \circ \varphi \circ \iota(a) = \tau \circ \varphi \circ \iota \circ \gamma_j(a), \quad j = 1, 2$$

for all $a \in C(X) \otimes \mathcal{U}$, where $\gamma_j = \alpha_j \otimes \sigma$ and σ is as in 2.6, $j = 1, 2$ and $\iota : C(X) \otimes \mathcal{U} \rightarrow C(X) \otimes \mathcal{U} \rtimes_{\gamma_1} \mathbb{Z}$ is the natural embedding. Let $\mathbb{D} = \rho_B(K_0(B))$. There is a unital simple AF-algebra B_1 with $(K_0(B_1), K_0(B_1)_+, [1_{B_1}]) = (\mathbb{D}, \mathbb{D}_+, 1)$ and there is a unital monomorphism $h_{00} : B \rightarrow B_1$ which maps $K_0(B)$ to $\mathbb{D}$ by the map ρ_B . Thus we may assume that $B = B_1$, without loss of generality. In particular, now we assume that $\ker \rho_B = \{0\}$. By replacing B by $B \otimes \mathcal{U}$, we may also assume that $K_0(B)$ is divisible.

Put $A_0 = C(X) \otimes \mathcal{U}$. Suppose that $e \in M_k(A_0 \rtimes_{\beta} \mathbb{Z})$ is a projection. Let $E : A_0 \rtimes_{\gamma_1} \mathbb{Z} \rightarrow A_0$ be the canonical conditional expectation. Since γ_1 has cyclic Rokhlin property, by Cor 4.2 of [32], there is a tracial state t of A_0 such that

$$t(E(a)) = \tau \circ \varphi(a) \quad (\text{e 8.247})$$

for all $a \in A_0 \rtimes_{\gamma_1} \mathbb{Z}$. Define $\bar{\gamma}_2 : A_0 \rtimes_{\gamma_1} \mathbb{Z} \rightarrow A_0 \rtimes_{\gamma_1} \mathbb{Z}$ by $\bar{\gamma}_2(a) = \gamma_2(a)$ for $a \in A_0$ and $\bar{\gamma}_2(u_{\gamma_1}) = u_{\gamma_1}$, where u_{γ_1} is the canonical unitary in $A_0 \rtimes_{\gamma_1} \mathbb{Z}$ implementing γ_1 . Since γ_1 commutes with γ_2 , $\bar{\gamma}_2$ defines an automorphism on $A_0 \rtimes_{\gamma_1} \mathbb{Z}$. For the canonical expectation $E : A_0 \rtimes_{\gamma_1} \mathbb{Z} \rightarrow A_0$, one computes that

$$E(\bar{\gamma}_2(a)) = \gamma_2(E(a)) \quad (\text{e 8.248})$$

for all $a \in A_0 \rtimes_{\gamma_1} \mathbb{Z}$.

Since $\tau \circ \varphi \circ \iota$ is γ_1 and γ_2 invariant, by applying (e 8.247) and (e 8.248)), we obtain that

$$t(E(\bar{\gamma}_2(a))) = t(\gamma_2(E(a))) = t(E(a)) \quad \text{for all } a \in A_0 \rtimes_{\gamma_1} \mathbb{Z}. \quad (\text{e 8.249})$$

Now let $e \in M_k(A_0 \rtimes_{\beta_1} \mathbb{Z})$ be a projection. It follows from (e 8.249) that

$$t(E(\bar{\gamma}_2(e))) = t(E(e)). \quad (\text{e 8.250})$$

It follows that

$$\tau \circ \varphi(\bar{\gamma}_2(e)) = \tau \circ \varphi(e). \quad (\text{e 8.251})$$

Since $\ker \rho_B = \{0\}$, we conclude that

$$\varphi_{*0} = (\varphi \circ \bar{\gamma}_2)_{*0} \quad (\text{e 8.252})$$

Since $K_1(B) = 0$ and $K_0(B)$ is divisible, we have

$$[\varphi] = [\varphi \circ \bar{\gamma}_2] \text{ in } KL(A_0 \rtimes_{\bar{\gamma}_1} \mathbb{Z}, B). \quad (\text{e 8.253})$$

It follows from 7.1, 7.3 and 7.6 that there exists a T -full (for some $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$) asymptotic embedding $\{\psi_n\}$ from $(A_0 \rtimes_{\gamma_1} \mathbb{Z}) \rtimes_{\bar{\gamma}_2} \mathbb{Z}$ into $B \otimes \mathcal{U} (\cong B)$ satisfying the following.

$$\lim_{n \rightarrow \infty} \|\psi_n(a) - h_n(a)\| = 0 \text{ for all } a \in A_0 \rtimes_{\gamma_1} \mathbb{Z}, \quad (\text{e 8.254})$$

where $h_n : A_0 \rtimes_{\gamma_1} \mathbb{Z} \rightarrow B \otimes \mathcal{U}$ is a unital monomorphism. Moreover, by 7.2, for each n ,

$$(\tau \otimes t) \circ h_n(a) = (\tau \otimes t) \circ h_1(a) \text{ for all } a \in A_0 \rtimes_{\gamma_1} \mathbb{Z}. \quad (\text{e 8.255})$$

Again,

$$[h_n] = [h_1] \text{ in } KL(A_0 \rtimes_{\gamma_1} \mathbb{Z}, B \otimes \mathcal{U}), \quad (\text{e 8.256})$$

$n = 1, 2, \dots$. It follows from (e 8.256), (e 8.255) and 3.6 that there exists a sequence of unitaries $\{w_n\} \subset B \otimes \mathcal{U}$ such that

$$\lim_{n \rightarrow \infty} \|\text{ad } w_n \circ h_n \circ j(f) - h_1 \circ j(f)\| = 0 \text{ for all } f \in C(X). \quad (\text{e 8.257})$$

Thus, (by also (e 8.254),

$$\lim_{n \rightarrow \infty} \|\text{ad } w_n \circ \psi_n \circ j_1(f) - \psi_1 \circ j_1(f)\| = 0 \quad (\text{e 8.258})$$

for all $f \in C(X)$. Now by replacing ψ_n by $\text{ad } w_n \circ \psi_n$, we may assume that

$$\lim_{n \rightarrow \infty} \|\psi_n \circ j(f) - h_1 \circ j(f)\| = 0 \text{ for all } f \in C(X). \quad (\text{e 8.259})$$

□

Corollary 8.2. *Let X be a compact metric space and let $\Lambda : \mathbb{Z}^2 \rightarrow \text{Aut}(C(X))$ be a $\mathbb{Z}^2$ -action on X . Suppose that there is a strictly positive Λ -invariant Borel probability measure μ . Then $C(X) \rtimes_{\Lambda} \mathbb{Z}^2$ is quasidiagonal.*

We would also mention the following two asymptotic embeddings:

Proposition 8.3. *Let A be a unital AH-algebra and let $\alpha \in \text{Aut}(A)$ be an automorphism. Then there is a T -full (for some $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}$) asymptotic embedding $\{\varphi_n\} : A \rtimes_{\alpha} \mathbb{Z} \rightarrow C$ for some unital simple AF-algebra C if and only if A admits a faithful α -invariant tracial state.*

Proof. By replacing A by $A \otimes \mathcal{U}$ and replacing α by $\alpha \otimes \sigma$, we may assume that α has the cyclic Rokhlin property of 2.6. By 4.2, there is a unital monomorphism $h : A \rightarrow B$ for some unital simple AF-algebra B with unique tracial state τ such that $\ker \rho_B = \{0\}$ such that

$$\tau \circ h = \tau \circ h \circ \alpha \quad (\text{e 8.260})$$

for all $\tau \in T(B)$. Since $\ker \rho_B = \{0\}$, it follows that

$$h_{*0} = (h \circ \alpha)_{*0} \quad (\text{e 8.261})$$

It follows from 7.5 that there is an asymptotic embedding $\{\varphi_n\} : A \rtimes_{\alpha} \mathbb{Z} \rightarrow B \otimes \mathcal{U}$ and a sequence of monomorphisms $h_n : A \rightarrow B \otimes \mathcal{U}$ such that

$$\lim_{n \rightarrow \infty} \|\varphi_n \circ j(a) - h_n(a)\| = 0 \text{ and}$$

$$\tau \circ h_n(a) = \tau \circ h_1(a) \text{ for all } a \in A.$$

Note since $K_1(B) = \{0\}$ and $K_0(B \otimes \mathcal{U})$ is divisible and $\ker \rho_B = \{0\}$,

$$[h_n] = [h_1] \text{ in } KL(A, B \otimes \mathcal{U}).$$

It follows from 3.6 that there is a sequence of unitaries $\{w_n\} \subset B$ such that

$$\lim_{n \rightarrow \infty} \|\text{ad } w_n \circ h_n(a) - h_1(a)\| = 0 \text{ for all } a \in A.$$

We now replace ψ_n by $\text{ad } w_n \circ \psi_n$. Thus, we may assume that

$$\lim_{n \rightarrow \infty} \|\psi_n \circ j(a) - h_1(a)\| = 0 \text{ for all } a \in A.$$

Since B is simple, it is easy to see that there is $T' : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$ such that $\{\psi_n \circ j\}$ is T' -full. Since we have assume that α has the cyclic Rokhlin property, by 7.6, $\{\psi_n\}$ is T -full for some $T : A_+ \setminus \{0\} \rightarrow \mathbb{N} \times \mathbb{R}_+$.

□

Proposition 8.4. *Let A be a separable amenable residually finite dimensional C^* -algebra in $\mathcal{N}$ and $\alpha \in \text{Aut}(A)$ such that $(\alpha)_{*0} = \text{id}|_{K_0(A)}$. Then there is an asymptotic embedding $\{\varphi_n\} : A \rtimes_{\alpha} \mathbb{Z} \rightarrow B$ for some unital simple AF-algebra.*

Proof. It follows from [25] that there is a unital monomorphism $h : A \rightarrow B$ for some unital simple AF-algebra B with a unique tracial state τ . By mapping B to another simple AF-algebra, if necessary, we may assume that $K_0(B)$ is divisible. Since $(\alpha)_{*0} = \text{id}|_{K_0(A)}$, it follows that

$$[h \circ \alpha] = [h] \text{ in } KL(A, B). \quad (\text{e8.262})$$

It follows from 7.4 that $A \rtimes_{\alpha} \mathbb{Z}$ can be asymptotically embedded into a unital simple AF-algebra B . □

References

- [1] B. Blackadar and E. Kirchberg *Generalized inductive limits of finite-dimensional C^* -algebras*, Math. Ann. **307** (1997), 343–380.
- [2] B. Blackadar and E. Kirchberg *Inner quasidiagonality and strong NF algebras*, Pacific J. Math. **198** (2001), 307–329.
- [3] L. G. Brown and M. Dădărlat, *Extensions of C^* -algebras and quasidiagonality*, J. London Math. Soc. **53** (1996), 582–600.
- [4] N. Brown, *AF embeddability of crossed products of AF algebras by the integers*, J. Funct. Anal. **160** (1998), no. 1, 150–175.
- [5] N. Brown, *Crossed products of UHF algebras by some amenable groups*, Hokkaido Math. J. **29** (2000), 201–211.
- [6] N. Brown, *On quasidiagonal C^* -algebras. Operator algebras and applications 19–64*, Adv. Stud. Pure Math., 38, Math. Soc. Japan, Tokyo, 2004.
- [7] N. Brown and M. Dadarlat, *Extensions of quasidiagonal C^* -algebras and K -theory*, Operator algebras and applications, 65–84, Adv. Stud. Pure Math., 38, Math. Soc. Japan, Tokyo, 2004.
- [8] M.D. Choi and E. Effros, *The completely positive lifting problem for C^* -algebras*, Ann. of Math. **104** (1976), 309–322.
- [9] M. Dădărlat, *Quasidiagonal morphisms and homotopy*, J. Funct. Anal. **151** (1997), 213–233.
- [10] M. Dădărlat, *On the approximation of quasidiagonal C^* -algebras*, J. Funct. Anal. **167** (1999), 69–78.
- [11] M. Dădărlat, *Residually finite dimensional C^* -algebras and subquotients of the CAR algebra*, Math. Res. Lett. **8** (2001), 545–555.
- [12] K. Davidson, D. Herrero and N. Salinas, *Quasidiagonal operators, approximation, and C^* -algebras*, Indiana Univ. Math. J. **38** (1989), 973–998.
- [13] S. Eilers, T. Loring and G. K. Pedersen, *Quasidiagonal extensions and AF algebras*, Math. Ann. **311** (1998), 233–249.

- [14] G. A. Elliott, *On the classification of C^* -algebras of real rank zero*, J. Reine Angew. Math. **443** (1993), 179–219.
- [15] G. A. Elliott and G. Gong, *On the classification of C^* -algebras of real rank zero. II*, Ann. of Math. **144** (1996), 497–610.
- [16] R. Exel, *The soft torus and applications to almost commuting matrices*, Pacific J. Math., **160** (1993), 207–217.
- [17] G. Gong and H. Lin, *Almost multiplicative morphisms and K -theory*, Internat. J. Math. **11** (2000), 983–1000.
- [18] D. Hadwin, *Strongly quasidiagonal C^* -algebras*, J. Operator Theory **18** (1987), 3–18.
- [19] S. Hu, H. Lin and Y. Xue, *The tracial topological rank of C^* -algebras. II*, Indiana Univ. Math. J. **53** (2004), 1578–1603.
- [20] A. Kishimoto, *Automorphisms of $\mathbf{AT}$ algebras with the Rohlin property*, J. Operator Theory **40** (1998), 277–294.
- [21] A. Kishimoto and A. Kumjian, *The Ext class of an approximately inner automorphism. II*, J. Operator Theory **46** (2001), 99–122.
- [22] L. Li, *Classification of simple C^* -algebras: inductive limits of matrix algebras over one-dimensional spaces*, J. Funct. Anal. **192** (2002), 1–51.
- [23] H. Lin, *Tracial topological ranks of C^* -algebras*, Proc. London Math. Soc., **83** (2001), 199–234.
- [24] H. Lin, *An introduction to the classification of amenable C^* -algebras*, World Scientific Publishing Co., Inc., River Edge, NJ, 2001. xii+320 pp. ISBN: 981-02-4680-3.
- [25] H. Lin, *Residually finite dimensional and AF-embeddable C^* -algebras*, Proc. Amer. Math. Soc. **129** (2001), 1689–1696.
- [26] H. Lin, *Classification of simple C^* -algebras and higher dimensional noncommutative tori*, Ann. of Math. (2) **157** (2003), no. 2, 521–544.
- [27] H. Lin, *Classification of simple C^* -algebras with tracial topological rank zero*, Duke Math. J., **125** (2004), 91–119.
- [28] H. Lin, *Traces and simple C^* -algebras with tracial topological rank zero*, J. Reine Angew. Math. **568** (2004), 99–137.
- [29] H. Lin, *An approximate universal coefficient theorem*, Trans. Amer. Math. Soc. **357** (2005), 3375–3405.
- [30] H. Lin, *The Rokhlin property for automorphisms on a unital simple C^* -algebra*, Contemp. Math., **414**, 2006, 189–215.
- [31] H. Lin, *Classification of homomorphisms and dynamical systems*, Trans. Amer. Math. Soc. **359** (2007), 859–895.
- [32] H. Lin, *Embedding crossed products into a unital simple AF-algebra*, preprint, arxiv.org/OA/0604047

- [33] H. Lin, *Approximate homotopy of homomorphisms from $C(X)$ into a simple C^* -algebra*, preprint, [arxiv.org/ OA/0612125](https://arxiv.org/abs/0612125).
- [34] H. Matui, *AF embeddability of crossed products of AT algebras by the integers and its application*, J. Funct. Anal. **192** (2002), 562–580.
- [35] C. Pasnicu and N. C. Phillips *Crossed products by $\mathbb{Z}$ with Rokhlin property*, in preparation.
- [36] N. C. Phillips, *Crossed products of the Cantor set by free minimal actions of $\mathbb{Z}^d$* , Comm. Math. Phys. **256** (2005), 1–42.
- [37] M. Pimsner, *Embedding some transformation group C^* -algebras into AF-algebras*, Ergodic Theory Dynam. Systems **3** (1983), 613–626.
- [38] M. Rørdam, *On the structure of simple C^* -algebras tensored with a UHF-algebra. II* J. Funct. Anal. **107** (1992), 255–269.
- [39] J. Rosenberg, *Appendix to Strongly quasidiagonal C^* -algebras*, J. Operator Theory **18** (1987), 3–18.
- [40] N. Salinas, *Relative quasidiagonality and KK -theory*, Houston J. Math., **18** (1992), 97–116.
- [41] F. J. Thayer, *Quasidiagonal C^* -algebras*, J. Funct. Anal. **25** (1977), 50–57.
- [42] J. Villadsen, *On the stable rank of simple C^* -algebras*, J. Amer. Math. Soc. **12** (1999) 1091–1102.
- [43] D. Voiculescu, *A non-commutative Weyl-von Neumann theorem*, Rev. Roumaine Math., Pures Apply. **21** (1976), 97–113.
- [44] D. Voiculescu, *Almost inductive limit automorphisms and embeddings into AF-algebras*, Ergodic Theory Dynam. Systems, **6** (1986), 475–484.
- [45] D. Voiculescu, *A note on quasi-diagonal C^* -algebras and homotopy*, Duke Math. J. **62** (1991), 267–271.
- [46] D. Voiculescu, *Around quasidiagonal operators* Integral Equations Operator Theory **17** (1993), 137–149.
- [47] S. Zhang, *K_1 -groups, quasidiagonality, and interpolation by multiplier projections*, Trans. Amer. Math. Soc. **325** (1991), 793–818.

Department of Mathematics
East China Normal University
Shanghai, China
and (current)
Department of Mathematics
University of Oregon
Eugene, Oregon 97403
U.S.A.